

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>7</sup> :</b> <b>C07D 231/44, 401/04, A01N 43/56</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 00/35884</b> <b>(43) International Publication Date:</b> 22 June 2000 (22.06.00)
<b>(21) International Application Number:</b> PCT/EP99/10452 <b>(22) International Filing Date:</b> 10 December 1999 (10.12.99)  <b>(30) Priority Data:</b> 60/111,857 11 December 1998 (11.12.98) US 60/140,680 24 June 1999 (24.06.99) US  <b>(71) Applicant:</b> AVENTIS CROPSCIENCES S.A. [FR/FR]; 55, avenue René Cassin, F-69009 Lyon (FR).  <b>(72) Inventors:</b> RIBEILL, Yves; 313 Meetinghouse Circle, Raleigh, NC 27609 (US). HUBER, Scot, Kevin; 6104 Bramblewood Drive, Raleigh, NC 27612 (US). MCCOMB, Susan, Marie; 110 Arrowhead Way, Cary, NC 27513 (US). MALASKA, Michael, James; 3228 Gait Way, Chapel Hill, NC 27516 (US). CHOU, David; 400 Dunwoody Drive, Raleigh, NC 27615 (US). PEREZ DE LEON, Adalberto; 716 Sarratt Ridge Court, Wake Forest, NC 27587 (US).  <b>(74) Agent:</b> AVENTIS CROPSCIENCE S.A.; Boîte postale 9163, F-69263 Lyon Cedex 9 (FR).		<b>(81) Designated States:</b> AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
<b>(54) Title:</b> CONTROL OF ARTHROPODS IN ANIMALS  <b>(57) Abstract</b>  A method of controlling parasites in or on an animal comprising administering to the animal a parasitically effective, substantially non-emetic 1-arylpyrazole.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakistan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

### Control of Arthropods in Animals

5           The present invention relates to a method of control of parasites in animals, compositions comprising a compound effective for the said control and new compounds effective against parasites.

10           It is generally a goal of agronomists and veterinarians to possess sufficient means to control pests, particularly arthropods, when they attempt to invade or attack mammals, particularly domestic animals and/or livestock. A classical method of controlling such pests has been the use of topical and/or systemic pesticides on or in the domestic animal which is being attacked. Generally effective treatments include the oral administration of insect growth regulators, such as lufenuron, or antihelminth compounds such as an ivermectin or an avermectin, or the  
15           topical application of the insecticide fipronil. It is advantageous to apply pesticides to animals in oral form so as to prevent the possible contamination of humans or the surrounding environment.


          It is an object of the present invention to provide new pesticides which may be used in domestic animals.

20           Another object of the invention is to provide safer pesticides for domestic animals.

          Another object of the invention is to provide new pesticides for domestic animals that may be used in lower doses than existing pesticides.

25           These objects are met in whole or in part by the present invention.

          The present invention provides a method of controlling parasites in or on an animal comprising administering orally to the animal a parasitically effective, substantially non-emetic amount of a 1-arylpyrazole of formula (I):



(I)

wherein:

R<sub>1</sub> is cyano, acetyl, C(S)NH<sub>2</sub>, alkyl, haloalkyl, C(=NOH)NH<sub>2</sub> or C(=NNH<sub>2</sub>)NH<sub>2</sub>;

5 R<sub>2</sub> is S(O)<sub>n</sub>R<sub>3</sub>; C<sub>2</sub>-C<sub>3</sub> alkenyl, C<sub>2</sub>-C<sub>3</sub> haloalkenyl, cycloalkyl, halocycloalkyl or C<sub>2</sub>-C<sub>3</sub> alkynyl;

R<sub>3</sub> is alkyl or haloalkyl;

R<sub>4</sub> is -N=C(R<sub>5</sub>)-Z-R<sub>6</sub>, -N=C(R<sub>5</sub>)-N(R<sub>7</sub>)-R<sub>8</sub>, or -N(R<sub>9</sub>)-C(R<sub>5</sub>)=NR<sub>6</sub>;

10 R<sub>5</sub> is hydrogen; alkyl; or alkyl substituted by halogen, alkoxy, haloalkoxy or -S(O)<sub>m</sub>R<sub>15</sub>;

R<sub>6</sub> and R<sub>7</sub> each independently represent hydrogen, alkyl, C<sub>3</sub>-C<sub>5</sub> alkenyl or C<sub>3</sub>-C<sub>5</sub> alkynyl; or

15 alkyl substituted by one or more halogen, alkoxy, haloalkoxy, amino, alkylamino, dialkylamino, cyano or -S(O)<sub>m</sub>R<sub>15</sub>; or alkyl substituted by phenyl or pyridyl each of which is optionally substituted with one or more groups selected from halogen, nitro and alkyl; or

20 R<sub>6</sub> and R<sub>7</sub> may form together with the nitrogen to which they are attached a 3 to 7 membered ring which may additionally contain one or more heteroatoms selected from oxygen, nitrogen or sulfur;

$R_8$  is alkoxy, haloalkoxy, amino, alkylamino, dialkylamino,  
 $R_{14}CO-$  or  $-S(O)_tR_{10}$ ;  
 $R_9$ ,  $R_{10}$  and  $R_{14}$  are alkyl or haloalkyl;  
 $R_{11}$  and  $R_{12}$  are independently selected from halogen, hydrogen,  
 5 CN and  $NO_2$ ;

$R_{13}$  is selected from halogen, haloalkyl, haloalkoxy,  $-S(O)_qCF_3$ ,  
 and  $-SF_5$ ;


$R_{15}$  is alkyl or haloalkyl;

$X$  is selected from nitrogen and  $C-R_{12}$ ;

10  $Z$  is O,  $S(O)_a$ ; or  $NR_7$ ;

$a$ ,  $m$ ,  $n$  and  $q$  are independently selected from 0, 1, and 2; and  
 $t$  is 0 or 2; and veterinarily acceptable salts thereof.

In another aspect, the present invention provides a method of  
 controlling parasites in or on an animal comprising administering orally  
 15 to the animal a parasitocidally effective, substantially non-emetic amount  
 of a 1-arylpyrazole of formula (XX):



(XX)

wherein:

20  $R_{201}$  is cyano,  $C(O)alkyl$ ,  $C(S)NH_2$ , alkyl,  $C(=NOH)NH_2$  or  
 $C(=NNH_2)NH_2$ ;

R<sub>202</sub> is S(O)<sub>h</sub>R<sub>203</sub>, C<sub>2</sub>-C<sub>3</sub> alkenyl, C<sub>2</sub>-C<sub>3</sub> haloalkenyl, cycloalkyl, halocycloalkyl or C<sub>2</sub>-C<sub>3</sub> alkynyl;

R<sub>203</sub> is alkyl or haloalkyl;

R<sub>204</sub> is -N(R<sub>205</sub>)C(O)CR<sub>206</sub>R<sub>207</sub>R<sub>208</sub>, -N(R<sub>205</sub>)C(O)aryl, or  
5 -N(R<sub>205</sub>)C(O)OR<sub>207</sub>;

R<sub>205</sub> is alkyl, haloalkyl, cycloalkyl, halocycloalkyl, cycloalkylalkyl, halocycloalkylalkyl, alkoxyalkyl, haloalkoxyalkyl, C<sub>3</sub>-C<sub>5</sub> alkenyl, C<sub>3</sub>-C<sub>5</sub> haloalkenyl, C<sub>3</sub>-C<sub>5</sub> alkynyl, C<sub>3</sub>-C<sub>5</sub> haloalkynyl;

R<sub>206</sub> is hydrogen, halogen, alkoxy, haloalkoxy, alkoxyalkyl, haloalkoxyalkyl, formyloxy, alkylcarbonyloxy, haloalkylcarbonyloxy, alkylthio, haloalkylthio, alkylsulfinyl, haloalkylsulfinyl, alkylsulfonyl, haloalkylsulfonyl, alkylamino, dialkylamino, haloalkylamino, di(haloalkyl)amino, cycloalkyloxy, halocycloalkyloxy, alkoxyalkoxy, haloalkoxyalkoxy, alkoxyalkoxyalkoxy, aryloxy, or arylalkoxy;

R<sub>207</sub> and R<sub>208</sub> are independently hydrogen, alkyl, haloalkyl, cycloalkyl, or halocycloalkyl; or R<sub>207</sub> and R<sub>208</sub> may form together with the carbon to which they are attached a 3 to 7 membered ring which additionally may contain one or more heteroatoms selected from nitrogen, oxygen and sulfur;

X<sub>1</sub> is selected from nitrogen and C-R<sub>212</sub>;

R<sub>211</sub> and R<sub>212</sub> are independently selected from halogen, hydrogen, CN and NO<sub>2</sub>;

R<sub>213</sub> is selected from halogen, haloalkyl, haloalkoxy, -S(O)<sub>k</sub>CF<sub>3</sub>, and -SF<sub>5</sub>; and

h and k are independently selected from 0, 1, and 2;  
and veterinarily acceptable salts thereof.

By the term "veterinarily acceptable salts" is meant salts the anions of which are known and accepted in the art for the formation of salts for veterinary use. Suitable acid addition salts, e.g. formed by

compounds of formulae (I) and (XX) containing a basic nitrogen atom, e.g. an amino group, include salts with inorganic acids, for example hydrochlorides, sulphates, phosphates and nitrates and salts with organic acids for example acetic acid.

5 Unless otherwise specified, alkyl and alkoxy groups are generally lower alkyl and alkoxy groups, that is having from one to six carbon atoms, preferably from one to four carbon atoms. Generally, the haloalkyl, haloalkoxy and alkylamino groups have from one to four carbon atoms. The haloalkyl and haloalkoxy groups can bear one or  
10 more halogen atoms; preferred groups of this type include  $-CF_3$  and  $-OCF_3$ . Cycloalkyl groups generally have from 3 to 6 carbon atoms, preferably from 3 to 5 carbon atoms and may be substituted by one or more halogen atoms. Alkenyl, haloalkenyl, alkynyl, and haloalkynyl groups generally contain from 3 to 5 carbon atoms. By the term aryl is  
15 generally meant phenyl, pyridyl, furyl, and thiophenyl, each of which is optionally substituted by one or more halogen, alkyl, haloalkyl, nitro, alkoxy, haloalkoxy, hydroxy, amino, alkylamino or dialkylamino. In compounds of formula (I), by the term substituted alkyl is meant alkyl which is substituted by one or more halogen, alkoxy, haloalkoxy, amino,  
20 alkylamino, dialkylamino, cyano or  $-S(O)_mR_{15}$ ; or alkyl substituted by phenyl or pyridyl each of which is optionally substituted with one or more groups selected from halogen, nitro and alkyl; wherein  $R_{15}$  is alkyl or haloalkyl and m is zero, one or two. Preferably in compounds of formula (I), alkyl groups are generally substituted by from one to five  
25 halogen atoms, preferably from one to three halogen atoms. Chlorine and fluorine atoms are preferred.

Compounds of formula (I) wherein  $R_4$  is  $-N=C(R_5)-Z-R_6$ , Z is  $NR_7$  and  $R_6$  represents a hydrogen atom may exist as the tautomeric

double bond isomer form  $\text{-NH-C(R}_5\text{)=N-R}_7$ . It is to be understood that both such forms are embraced by the present invention.

In compounds of formula (XX) the following examples of radicals are provided:

- 5           An example of cycloalkylalkyl is cyclopropylmethyl;  
          an example of cycloalkoxy is cyclopropyloxy;  
          an example of alkoxyalkyl is  $\text{CH}_3\text{OCH}_2\text{-}$ ;  
          an example of alkoxyalkoxy is  $\text{CH}_3\text{OCH}_2\text{O-}$ ;  
          An example of alkoxyalkoxyalkoxy is  $\text{CH}_3\text{OCH}_2\text{OCH}_2\text{O-}$ ;  
10          An example of aryloxy is the phenoxy radical; and  
          An example of the arylalkoxy radical is benzyloxy or 2-phenylethoxy.

          Generally, in dialkylamino or di(haloalkyl)amino radicals, the alkyl and haloalkyl groups on nitrogen may be chosen independently of  
15          one another.

          It is also to be understood that enantiomeric and diastereomeric forms of the compounds of formulae (I) and (XX) and salts thereof are embraced by the present invention. Compounds of formula (I) may be generally prepared according to known processes, for example as  
20          described in European Patent Application 511845 or other processes according to the knowledge of a man skilled in the art of chemical synthesis.

          By the term non-emetic is meant a compound that does not generally elicit emesis from the animal when a protective, preventative  
25          or cleaning dose is administered to the animal. By the term emesis is meant vomiting. Generally an emetic substance elicits the said emesis in less than 24 hours after administration, preferably less than 8 hours, more preferably less than 2 hours. Generally when the compounds of the invention are administered to a population of animals, more than



70% of the animals are free of emesis, preferably more than 80%, most preferably more than 90%.

A preferred class of compounds of formula (I) for use in the control of parasites in animals are those wherein:

- 5            $R_1$  is cyano or alkyl;  
             $R_2$  is  $S(O)_nR_3$ ;  
             $R_3$  is alkyl or haloalkyl;  
             $R_4$  is  $-N=C(R_5)-Z-R_6$ ;  
             $R_5$  is hydrogen, alkyl or haloalkyl;  
10            $Z$  is O,  $S(O)_a$ ; or  $NR_7$ ;  
             $R_6$  and  $R_7$  are independently selected from hydrogen and unsubstituted or substituted alkyl; or  
             $R_6$  and  $R_7$  may form together with the nitrogen to which they are attached a 3 to 7 membered ring which may additionally contain one or  
15           more heteroatoms selected from oxygen, nitrogen or sulfur;  $X$  is selected from nitrogen and C- $R_{12}$ ;  
             $R_{11}$  and  $R_{12}$  are independently selected from halogen, hydrogen, CN and  $NO_2$ ;  
             $R_{13}$  is selected from halogen, haloalkyl, haloalkoxy,  $-S(O)_qCF_3$ ,  
20           and  $-SF_5$ ;  
            a, n and q are independently selected from 0, 1, and 2.

- 3)           Preferably  $R_6$  is alkyl which is substituted by one or more halogen, alkoxy, haloalkoxy, amino, alkylamino, dialkylamino, sulfide,  
25           sulfoxide, sulfone, or phenyl or pyridyl moieties of which each phenyl or pyridyl moiety is optionally substituted with one or more groups selected from halo, nitro, and alkyl.

Preferably the method of the invention has one or more of the following features:

$R_1$  is cyano;

$R_4$  is  $-N=C(R_5)-Z-R_6$  and  $Z$  is  $-NR_7$ ;

$X$  is  $C-R_{12}$ ;  $R_{11}$  and  $R_{12}$  represent a chlorine atom; and  $R_{13}$  is  $CF_3$ ,  $OCF_3$  or  $-SF_5$ ;

5  $R_{12}$  is  $-S(O)_nCF_3$  and  $n$  is 0, 1, or 2.

A further preferred class of compounds of formula (I) for use in the control of parasites in animals are those wherein:

10  $R_1$  is cyano or alkyl;  $R_4$  is  $-N=C(R_5)-Z-R_6$ ; and  $R_5$  is hydrogen or  $C_1-C_3$  alkyl.

The compounds of formula (I) for use in the control of parasites in animals, preferably have one or more of the following features:

$R_1$  is cyano or methyl;

$R_3$  is halomethyl (preferably  $CF_3$ );

15  $R_{11}$  and  $R_{12}$  each independently represent a halogen atom;

$X$  is  $C-R_{12}$ ;

$R_{13}$  is haloalkyl (preferably  $CF_3$ ), haloalkoxy (preferably  $OCF_3$ ), or  $-SF_5$ ; or

$n$  is 0, 1 or 2 (preferably 0 or 1).

20 A further preferred class of compounds of formula (I) for use in the control of parasites in animals are those wherein:

$R_1$  is cyano;

$R_2$  is  $S(O)_nR_3$ ;

$R_3$  is halomethyl;

25  $R_4$  is  $-N=C(R_5)-Z-R_6$ ;

$Z$  is  $NR_7$ ;

$R_5$  is hydrogen or alkyl;

R<sub>6</sub> and R<sub>7</sub> each independently represent hydrogen, alkyl, alkenyl or alkynyl; or alkyl substituted by one or more halogen, alkoxy, haloalkoxy, amino, alkylamino, dialkylamino, cyano or -S(O)<sub>m</sub>R<sub>15</sub>; or alkyl substituted by phenyl or pyridyl which rings are optionally substituted with one or more groups selected from halogen, nitro and alkyl;

X is selected from nitrogen and C-R<sub>12</sub>;

R<sub>11</sub> and R<sub>12</sub> each independently represent a halogen atom;

R<sub>13</sub> is selected from haloalkyl, haloalkoxy and -SF<sub>5</sub>;

R<sub>15</sub> is alkyl or haloalkyl; and

m and n are independently selected from 0, 1, and 2.

A further preferred class of compounds of formula (I) for use in the control of parasites in animals is that wherein:

R<sub>1</sub> is cyano;

R<sub>2</sub> is S(O)<sub>n</sub>CF<sub>3</sub>;

R<sub>4</sub> is -N=C(R<sub>5</sub>)-Z-R<sub>6</sub> or -N=C(R<sub>5</sub>)-N(R<sub>7</sub>)-R<sub>8</sub>;

Z is NR<sub>7</sub>;

R<sub>5</sub> is hydrogen or alkyl;

R<sub>6</sub> and R<sub>7</sub> each independently represent hydrogen, alkyl, alkenyl or alkynyl; or alkyl substituted by one or more halogen, alkoxy, haloalkoxy, amino, alkylamino, dialkylamino, cyano or -S(O)<sub>m</sub>R<sub>15</sub>; or methyl substituted by phenyl or pyridyl which rings are optionally substituted with one or more groups selected from halogen, nitro and alkyl;

R<sub>8</sub> is alkoxy, haloalkoxy, amino, alkylamino, dialkylamino or -S(O)<sub>t</sub>R<sub>10</sub>;

X is selected from nitrogen and C-R<sub>12</sub>;

R<sub>10</sub> and R<sub>15</sub> independently represent alkyl or haloalkyl;

$R_{11}$  and  $R_{12}$  each represent a chlorine atom;

$R_{13}$  is  $CF_3$  or  $-SF_5$ ; and

$m$  and  $n$  are 0, 1 or 2; and  $t$  is 0 or 2.

5 A further preferred class of compounds of formula (I) for use in the control of parasites in animals are those wherein:

$R_1$  is cyano;

$R_2$  is  $S(O)_nCF_3$ ;

$R_4$  is  $-N=C(R_5)-Z-R_6$ ;

$Z$  is  $NR_7$ ;

10  $R_5$  is hydrogen or methyl;

$R_6$  and  $R_7$  each independently represent hydrogen, alkyl, alkenyl or alkynyl; or alkyl substituted by one or more halogen, alkoxy, haloalkoxy, amino, alkylamino, dialkylamino, cyano or  $-S(O)_mR_{15}$ ; or alkyl substituted by phenyl or pyridyl which rings are optionally substituted with one or more groups selected from halogen, nitro and

15 alkyl;

$X$  is  $C-R_{12}$ ;

$R_{11}$  and  $R_{12}$  each represent a chlorine atom;

$R_{13}$  is  $CF_3$  or  $-SF_5$ ;

20  $R_{15}$  is alkyl or haloalkyl;

$m$  is zero, one or two; and

$n$  is 0 or 1.

A further preferred class of compounds of formula (I) for use in the control of parasites in animals are those wherein:

25  $R_1$  is cyano;

$R_2$  is  $S(O)_nCF_3$ ;

$R_4$  is  $-N=C(R_5)-Z-R_6$ ;

$Z$  is  $NR_7$ ;

R<sub>5</sub> and R<sub>7</sub> each represent a hydrogen atom;

R<sub>6</sub> is alkyl or haloalkyl;

X is C-R<sub>12</sub>;

R<sub>11</sub> and R<sub>12</sub> each represent a chlorine atom;

5 R<sub>13</sub> is CF<sub>3</sub> or -SF<sub>5</sub>; and

n is 0.

Compounds of formula (XX) which are preferred according to the present invention are those wherein:

10 wherein:

R<sub>201</sub> is cyano;

R<sub>202</sub> is S(O)<sub>h</sub>R<sub>203</sub>;

R<sub>203</sub> is alkyl or haloalkyl;

R<sub>204</sub> is -N(R<sub>205</sub>)C(O)CR<sub>206</sub>R<sub>207</sub>R<sub>208</sub>;

15 R<sub>205</sub> is alkyl, haloalkyl, cycloalkyl, cycloalkylalkyl and halocycloalkylalkyl;

R<sub>206</sub> is alkoxy, haloalkoxy, or hydrogen;

R<sub>207</sub> and R<sub>208</sub> are independently hydrogen, alkyl, or haloalkyl; or

20 R<sub>207</sub> and R<sub>208</sub> may form together with the carbon to which they are attached a 3 to 7 membered ring which additionally may contain one or more heteroatoms selected from nitrogen, oxygen and sulfur;

X<sub>1</sub> is selected from nitrogen and C-R<sub>212</sub>;

R<sub>211</sub> and R<sub>212</sub> are independently selected from halogen, hydrogen, CN and NO<sub>2</sub>;

25 R<sub>213</sub> is selected from halogen, haloalkyl, haloalkoxy, -S(O)<sub>k</sub>CF<sub>3</sub>, and -SF<sub>5</sub>; and

h and k are independently selected from 0, 1, and 2.

A preferred group of compounds of formula (XX) is that wherein the ring which is formed by R<sub>207</sub> and R<sub>208</sub> is interrupted by one or more heteroatoms, more preferably one oxygen atom.

5 The compounds of formula (I) of the present invention preferably have one or more of the following features:

R<sub>201</sub> is cyano;

R<sub>203</sub> is halomethyl, preferably CF<sub>3</sub>;

R<sub>211</sub> and R<sub>212</sub> are independently halogen;

X<sub>1</sub> is C-R<sub>212</sub>;

10 R<sub>213</sub> is haloalkyl, haloalkoxy or -SF<sub>5</sub>; or

h is 0 or 1, or 2, preferably 0 or 1.

A preferred class of compounds that wherein R<sub>204</sub> is  
N(R<sub>205</sub>)C(O)CR<sub>206</sub>R<sub>207</sub>R<sub>208</sub>.

15 Another preferred class of compounds that wherein R<sub>204</sub> is  
N(R<sub>205</sub>)C(O)aryl.


Another preferred class of compounds that wherein R<sub>204</sub> is  
N(R<sub>205</sub>)C(O)OR<sub>207</sub>.

Preferably R<sub>205</sub> is C<sub>1</sub>-C<sub>4</sub> alkyl, more preferably C<sub>1</sub>-C<sub>2</sub> alkyl, most preferably methyl.

20 Preferably R<sub>206</sub> is alkoxy, most preferably methoxy, ethoxy or propoxy.

Preferably R<sub>207</sub> and R<sub>208</sub> are both hydrogen.

25 Among the compounds which may be used in the invention some are new and hence in another aspect of the present invention there is provided a compound of formula (II):



(II)

wherein:

R<sub>21</sub> is cyano, alkyl, haloalkyl, , acetyl or -C(=S)NH<sub>2</sub>,

C(=NOH)NH<sub>2</sub> or C(=NNH<sub>2</sub>)NH<sub>2</sub>;

5

R<sub>22</sub> is S(O)<sub>m</sub>R<sub>23</sub>;

R<sub>23</sub> is alkyl or haloalkyl;

R<sub>24</sub> is -N=C(R<sub>25</sub>)N(R<sub>26</sub>)(R<sub>27</sub>) or -N=C(R<sub>25</sub>)-N(R<sub>27</sub>)-R<sub>28</sub>;

R<sub>25</sub> represents hydrogen or alkyl; or alkyl substituted by one or more halogen, alkoxy, haloalkoxy or -S(O)<sub>m</sub>R<sub>35</sub>;

10

R<sub>26</sub> and R<sub>27</sub> each independently represent hydrogen, alkyl, alkenyl or alkynyl; or alkyl substituted by one or more halogen, alkoxy, haloalkoxy, amino, alkylamino, dialkylamino, cyano or -S(O)<sub>m</sub>R<sub>35</sub>; or alkyl substituted by phenyl or pyridyl which rings are optionally substituted with one or more groups selected from halogen, nitro and alkyl; wherein R<sub>35</sub> is alkyl or haloalkyl and m is zero, one or two;

15

X is selected from nitrogen and C-R<sub>32</sub>;

R<sub>28</sub> is alkoxy, haloalkoxy, amino, alkylamino, dialkylamino or -S(O)<sub>t</sub>R<sub>30</sub>;

R<sub>30</sub> is alkyl or haloalkyl;

20

R<sub>31</sub> and R<sub>32</sub> are independently selected from halogen, hydrogen, CN and NO<sub>2</sub>;

$R_{33}$  is selected from halogen, haloalkyl, haloalkoxy,  $-S(O)_rCF_3$ , and  $-SF_5$ ;

m and r are independently selected from 0, 1, and 2; and t is 0 or 2; with the exclusion of the compound wherein  $R_{21}$  is cyano;  $R_{22}$  is  $-SCF_2CH_3$ ;  $R_{25}$  is hydrogen; X is  $C-R_{32}$ ;  $R_{26}$  and  $R_{27}$  are methyl;  $R_{31}$  and  $R_{32}$  are chlorine; and  $R_{33}$  is trifluoromethyl; and veterinarily acceptable salts thereof.

A further class of novel compounds of formula (II) are those wherein:

$R_{21}$  is cyano or methyl;

$R_{22}$  is  $S(O)_mR_{23}$ ;

$R_{23}$  is haloalkyl;

$R_{24}$  is  $-N=C(R_{25})N(R_{26})(R_{27})$ ;

$R_{25}$  and  $R_{27}$  are hydrogen or unsubstituted or substituted alkyl;

$R_{26}$  is haloalkyl;

X is selected from nitrogen and  $C-R_{32}$ ;

$R_{31}$  and  $R_{32}$  are independently selected from halogen, hydrogen, CN and  $NO_2$ ;

$R_{33}$  is selected from halogen, haloalkyl, haloalkoxy,  $-S(O)_rCF_3$ , and  $-SF_5$ ;

m and r are independently selected from 0, 1, and 2; with the exclusion of the compound wherein  $R_{21}$  is cyano;  $R_{22}$  is  $-SCF_2CH_3$ ;  $R_{25}$  is hydrogen; X is  $C-R_{32}$ ;  $R_{26}$  and  $R_{27}$  are methyl;  $R_{31}$  and  $R_{32}$  are chlorine; and  $R_{33}$  is trifluoromethyl.

A preferred class of novel compounds of formula (II) are those wherein:

$R_{21}$  is cyano;

$R_{22}$  is  $S(O)_mR_{23}$ ;



R<sub>23</sub> is halomethyl;

R<sub>24</sub> is -N=C(R<sub>25</sub>)N(R<sub>26</sub>)(R<sub>27</sub>);

R<sub>25</sub> is hydrogen or alkyl;

R<sub>26</sub> and R<sub>27</sub> each independently represent hydrogen, alkyl,

5 alkenyl or alkynyl; or alkyl substituted by one or more halogen, alkoxy, haloalkoxy, amino, alkylamino, dialkylamino, cyano or -S(O)<sub>m</sub>R<sub>15</sub>; or alkyl substituted by phenyl or pyridyl each of which is optionally substituted with one or more groups selected from halogen, nitro and alkyl; wherein R<sub>15</sub> is alkyl or haloalkyl and m is zero, one or two;

10 X is selected from nitrogen and C-R<sub>32</sub>;

R<sub>31</sub> and R<sub>32</sub> each represent a chlorine atom;

R<sub>33</sub> is selected from haloalkyl, haloalkoxy and -SF<sub>5</sub>;

m is selected from 0, 1, and 2.

A further preferred class of novel compounds of formula (II) are  
15 those wherein:

R<sub>21</sub> is cyano;

R<sub>22</sub> is S(O)<sub>m</sub>CF<sub>3</sub>;

R<sub>24</sub> is -N=C(R<sub>25</sub>)N(R<sub>26</sub>)(R<sub>27</sub>); or -N=C(R<sub>25</sub>)-N(R<sub>27</sub>)-R<sub>28</sub>;

R<sub>25</sub> is hydrogen or methyl;

20 R<sub>26</sub> and R<sub>27</sub> each independently represent hydrogen, alkyl, alkenyl or alkynyl; or alkyl substituted by one or more halogen, alkoxy, haloalkoxy, amino, alkylamino, dialkylamino, cyano or -S(O)<sub>m</sub>R<sub>15</sub>; or methyl substituted by phenyl or pyridyl which rings are optionally substituted with one or more groups selected from halogen, nitro and  
25 alkyl; wherein R<sub>15</sub> is alkyl or haloalkyl and m is zero, one or two;

R<sub>28</sub> is alkoxy, haloalkoxy, amino, alkylamino, dialkylamino or -S(O)<sub>t</sub>R<sub>30</sub>;

X is selected from nitrogen and C-R<sub>32</sub>;

R<sub>30</sub> is alkyl or haloalkyl;

R<sub>31</sub> and R<sub>32</sub> each represent a chlorine atom;

R<sub>33</sub> is CF<sub>3</sub> or -SF<sub>5</sub>; and

m is 0, 1 or 2; and t is 0 or 2.

5 A more preferred class of novel compounds of formula (II) are those wherein:

R<sub>21</sub> is cyano;

R<sub>22</sub> is S(O)<sub>m</sub>CF<sub>3</sub>;

R<sub>24</sub> is -N=C(R<sub>25</sub>)N(R<sub>26</sub>)(R<sub>27</sub>);

10 R<sub>25</sub> and R<sub>27</sub> each independently represent hydrogen or methyl;

R<sub>26</sub> represents hydrogen, alkyl, alkenyl or alkynyl; or alkyl substituted by one or more halogen, alkoxy, haloalkoxy, amino, alkylamino, dialkylamino, cyano or -S(O)<sub>m</sub>R<sub>15</sub>; or methyl substituted by phenyl or pyridyl which rings are optionally substituted with one or more groups selected from halogen, nitro and alkyl; wherein R<sub>15</sub> is  
15 alkyl or haloalkyl and m is zero, one or two;

X is selected from nitrogen and C-R<sub>32</sub>;

R<sub>31</sub> and R<sub>32</sub> each represent a chlorine atom;

R<sub>33</sub> is CF<sub>3</sub> or -SF<sub>5</sub>; and

20 m is 0, 1 or 2.

An especially preferred class of novel compounds of formula (II) are those wherein:

R<sub>21</sub> is cyano;

R<sub>22</sub> is S(O)<sub>m</sub>CF<sub>3</sub>;

25 R<sub>24</sub> is -N=C(R<sub>25</sub>)N(R<sub>26</sub>)(R<sub>27</sub>);

R<sub>25</sub> and R<sub>27</sub> each represent a hydrogen atom;

R<sub>26</sub> is alkyl or (preferably) haloalkyl;

X is C-R<sub>32</sub>;

$R_{31}$  and  $R_{32}$  each represent a chlorine atom;

$R_{33}$  is  $CF_3$  or  $-SF_5$ ; and

$m$  is 0.

5 In another aspect of the present invention there is provided an compound of formula (XX) or a salt thereof as hereinbefore defined, provided that the compound is not 3-cyano-1-(2,6-dichloro-4-trifluoromethylphenyl)-5-(N-ethoxycarbonyl-N-methyl)amino-4-trifluoromethylthiopyrazole.

10

Most preferably, the following compounds of formula (I) and (XX) are preferred according to the present invention as listed in Tables 1 to 13. The Compound Numbers are for identification purposes only. The following symbols are hereby defined: Me means methyl; Et means  
15 ethyl; n-Pr means n-propyl; i-Pr means isopropyl; n-Bu means n-Butyl, and n-Pent means n-Pentyl; Cy means cyclopropyl.

Table 1

Compounds of formula (I) wherein  $R_1$  is cyano;  $R_2$  is  $SCF_3$ ;  $R_{11}$  is Cl, X is C-Cl,  $R_4$  is  $-N=C(R_5)ZR_6$ , Z is  $NR_7$ ,  $R_7$  is H, and  $R_{13}$  is  
20  $CF_3$  or  $SF_5$ .

Compound Number $R_{13}=CF_3$	Compound Number $R_{13}=SF_5$	$R_5$	$R_6$
201-1	201-2	Me	Me
202-1	202-2	Me	Et
203-1	203-2	Me	n-Pr
204-1	204-2	Me	i-Pr
205-1	205-2	Me	n-Bu
206-1	206-2	H	H

207-1	207-2	H	Et
208-1	208-2	H	n-Pr
209-1	209-2	H	i-Pr
210-1	210-2	H	n-Bu
211-1	211-2	H	CH <sub>2</sub> CF <sub>3</sub>
212-1	212-2	H	(CH <sub>2</sub> ) <sub>2</sub> CF <sub>3</sub>
213-1	213-2	H	CH <sub>2</sub> OMe
214-1	214-2	H	(CH <sub>2</sub> ) <sub>2</sub> OCF <sub>3</sub>
215-1	215-2	Me	CH <sub>2</sub> CF <sub>3</sub>
216-1	216-2	Me	(CH <sub>2</sub> ) <sub>2</sub> CF <sub>3</sub>
217-1	217-2	Me	(CH <sub>2</sub> ) <sub>2</sub> OMe
218-1	218-2	Me	(CH <sub>2</sub> ) <sub>2</sub> NMe <sub>2</sub>

**Table 2**

Compounds of formula (I) wherein R<sub>1</sub> is cyano; R<sub>11</sub> is Cl; R<sub>4</sub> is -  
N=C(R<sub>5</sub>)ZR<sub>6</sub> and Z is NR<sub>7</sub>.

Cmp d No.	R2	R5	R6	R7	X	R13
219	SCF <sub>3</sub>	H	CH <sub>3</sub>	CH <sub>3</sub>	C-Cl	CF <sub>3</sub>
220	SO <sub>2</sub> CF <sub>3</sub>	H	CH <sub>3</sub>	CH <sub>3</sub>	C-Cl	CF <sub>3</sub>
221	SCF <sub>3</sub>	H	CH <sub>2</sub> CN	H	C-Cl	CF <sub>3</sub>
222	SCF <sub>3</sub>	H	CH <sub>3</sub>	H	C-Cl	CF <sub>3</sub>
223	SOCF <sub>3</sub>	H	CH <sub>2</sub> Ph	H	C-Cl	CF <sub>3</sub>
224	SCF <sub>3</sub>	H	CH <sub>2</sub> Ph	H	C-Cl	CF <sub>3</sub>
225	SOCF <sub>3</sub>	H	CH <sub>3</sub>	H	C-Cl	CF <sub>3</sub>
226	SOCF <sub>3</sub>	H	CH <sub>3</sub>	H	C-Cl	CF <sub>3</sub>
227	SOCF <sub>3</sub>	H	CH <sub>2</sub> CF <sub>3</sub>	H	C-Cl	CF <sub>3</sub>
228	SO <sub>2</sub> CF <sub>3</sub>	H	2-propynyl	H	C-Cl	CF <sub>3</sub>
229	SO <sub>2</sub> CF <sub>3</sub>	H	CH <sub>2</sub> Ph	H	C-Cl	CF <sub>3</sub>
230	SO <sub>2</sub> CF <sub>3</sub>	H	CH <sub>2</sub> CF <sub>3</sub>	H	C-Cl	CF <sub>3</sub>
231	SOCF <sub>3</sub>	H	CH <sub>3</sub>	CH <sub>3</sub>	C-Cl	CF <sub>3</sub>
232	SCF <sub>3</sub>	H	CH <sub>2</sub> CF <sub>3</sub>	H	C-Cl	CF <sub>3</sub>
233	SCF <sub>3</sub>	H	CH <sub>2</sub> CF <sub>3</sub>	H	C-Cl	CF <sub>3</sub>
234	SCF <sub>3</sub>	H	2-propenyl	H	C-Cl	CF <sub>3</sub>
235	SCF <sub>3</sub>	H	2-propynyl	H	C-Cl	CF <sub>3</sub>

236	SOCF3	H	2-propynyl	H	C-Cl	CF3
237	SCF3	H	CH2OEt	H	N	CF3
238	SCF3	H	CH2OCH2CF3	CH3	C-Cl	CF3
239	SO2CF3	H	CH2CF3	CH3	C-Cl	SF5
240	SCF3	H	CH2CF3	H	N	SF5
241	SOCF3	H	CH2CH2CF3	H	C-Cl	CF3
242	SO2CF3	H	(CH2)3CF3	H	C-Cl	CF3
243	SCF3	H	(CH2)2N(CH3)2	H	C-Cl	CF3
244	SO2CF3	CH3	CH2(4-Cl Ph)	H	C-Cl	CF3
245	SCF3	H	CH2SO2CF3	H	C-Cl	CF3
246	SCF3	H	CH2(4-pyridyl)	H	C-Cl	CF3
247	SCF3	H	CH2(3-NO2 Ph)	H	C-Cl	CF3
248	SCF3	H	CH2CH2SCH3	H	C-Cl	CF3
249	SOCF3	H	CH2CF3	CH3	C-Cl	CF3

Note: Compound number 232 is the acetate salt, and compound number 233 is the citrate salt.

**Table 3**

5

Compounds of formula (I) wherein R<sub>1</sub> is cyano; R<sub>11</sub> is Cl; and R<sub>4</sub> is -N=C(R<sub>5</sub>)-N(R<sub>7</sub>)-R<sub>8</sub>.

Cmpd No.	R2	R5	R8	R7	X	R13
250	SCF3	H	OEt	H	C-Cl	CF3
251	SCF3	H	NHCH3	H	C-Cl	CF3
252	SCF3	H	NHCH3	CH3	C-Cl	CF3
253	SCF3	H	OCH2CF3	H	C-Cl	CF3
254	SCF3	H	N(CH3)2	H	N	SF5
255	SCF3	H	NH2	H	C-Cl	CF3
256	SCF3	H	S-nPr	H	C-Cl	CF3
257	SO2CF3	Et	S-nPr	H	C-Cl	SF5
258	SCF3	H	SO2CH3	H	C-Cl	CF3

The following compounds of formula (XX) are preferred according to the present invention as listed in Tables 4-12.

10

**Table 4**

Compounds of formula (XX) wherein R<sub>201</sub> is cyano; R<sub>202</sub> is SCF<sub>3</sub>; R<sub>204</sub> is N(R<sub>205</sub>)C(O)CR<sub>206</sub>R<sub>207</sub>R<sub>208</sub>; R<sub>207</sub> and R<sub>208</sub> =H; R<sub>211</sub> is Cl, X<sub>1</sub> is C-Cl, and R<sub>213</sub> is CF<sub>3</sub> or SF<sub>5</sub>.

Compound Number (R <sub>213</sub> =CF <sub>3</sub> )	Compound Number (R <sub>213</sub> =SF <sub>5</sub> )	R <sub>205</sub>	R <sub>206</sub>
1-1	1-2	Me	H
2-1	2-2	Me	OMe
3-1	3-2	Me	OEt
4-1	4-2	Me	O-i-Pr
5-1	5-2	Me	O-n-Bu
6-1	6-2	Et	H
7-1	7-2	Et	OMe
8-1	8-2	Et	OEt
9-1	9-2	Et	O-i-Pr
10-1	10-2	Et	O-n-Bu
11-1	11-2	n-Pr	H
12-1	12-2	n-Pr	OMe
13-1	13-2	n-Pr	OEt
14-1	14-2	n-Pr	O-i-Pr
15-1	15-2	n-Pr	O-n-Bu
16-1	16-2	i-Pr	H
17-1	17-2	i-Pr	OMe
18-1	18-2	i-Pr	OEt
19-1	19-2	i-Pr	O-i-Pr
20-1	20-2	i-Pr	O-n-Bu
21-1	21-2	n-Bu	H
22-1	22-2	n-Bu	OMe
23-1	23-2	n-Bu	OEt
24-1	24-2	n-Bu	O-i-Pr
25-1	25-2	n-Bu	O-n-Bu
26-1	26-2	CH <sub>2</sub> Cy	H
27-1	27-2	CH <sub>2</sub> Cy	OMe
28-1	28-2	CH <sub>2</sub> Cy	OEt

29-1	29-2	CH <sub>2</sub> Cy	O-i-Pr
30-1	30-2	CH <sub>2</sub> Cy	O-n-Bu
31-1	31-2	CH <sub>2</sub> CCH	H
32-1	32-2	CH <sub>2</sub> CCH	OMe
33-1	33-2	CH <sub>2</sub> CCH	OEt
34-1	34-2	CH <sub>2</sub> CCH	O-i-Pr
35-1	35-2	CH <sub>2</sub> CCH	O-n-Bu
36-1	36-2	Me	OAc
37-1	37-2	Me	CH <sub>2</sub> OMe
38-1	38-2	Me	CH <sub>2</sub> OEt
39-1	39-2	Me	O-i-Pr
40-1	40-2	Me	O-n-Bu
41-1	41-2	Me	OCH <sub>2</sub> CF <sub>3</sub>

Table 5

Compounds of formula (XX) wherein R<sub>201</sub> is cyano; R<sub>202</sub> is

S(O)CF<sub>3</sub>; R<sub>204</sub> is N(R<sub>205</sub>)C(O)CR<sub>206</sub>R<sub>207</sub>R<sub>208</sub>; R<sub>207</sub> and R<sub>208</sub> =H;

R<sub>211</sub> is Cl, X<sub>1</sub> is C-Cl, and R<sub>213</sub> is CF<sub>3</sub> or SF<sub>5</sub>.

5

Compound Number (R <sub>213</sub> =CF <sub>3</sub> )	Compound Number (R <sub>213</sub> =SF <sub>5</sub> )	R <sub>205</sub>	R <sub>206</sub>
1-3	1-4	Me	H
2-3	2-4	Me	OMe
3-3	3-4	Me	OEt
4-3	4-4	Me	O-i-Pr
5-3	5-4	Me	O-n-Bu
6-3	6-4	Et	H
7-3	7-4	Et	OMe
8-3	8-4	Et	OEt
9-3	9-4	Et	O-i-Pr
10-3	10-4	Et	O-n-Bu

11-3	11-4	n-Pr	H
12-3	12-4	n-Pr	OMe
13-3	13-4	n-Pr	OEt
14-3	14-4	n-Pr	O-i-Pr
15-3	15-4	n-Pr	O-n-Bu
16-3	16-4	i-Pr	H
17-3	17-4	i-Pr	OMe
18-3	18-4	i-Pr	OEt
19-3	19-4	i-Pr	O-i-Pr
20-3	20-4	i-Pr	O-n-Bu
21-3	21-4	n-Bu	H
22-3	22-4	n-Bu	OMe
23-3	23-4	n-Bu	OEt
24-3	24-4	n-Bu	O-i-Pr
25-3	25-4	n-Bu	O-n-Bu
26-3	26-4	CH <sub>2</sub> Cy	H
27-3	27-4	CH <sub>2</sub> Cy	OMe
28-3	28-4	CH <sub>2</sub> Cy	OEt
29-3	29-4	CH <sub>2</sub> Cy	O-i-Pr
30-3	30-4	CH <sub>2</sub> Cy	O-n-Bu
31-3	31-4	CH <sub>2</sub> CCH	H
32-3	32-4	CH <sub>2</sub> CCH	OMe
33-3	33-4	CH <sub>2</sub> CCH	OEt
34-3	34-4	CH <sub>2</sub> CCH	O-i-Pr
35-3	35-4	CH <sub>2</sub> CCH	O-n-Bu
36-3	36-4	Me	OAc
37-3	37-4	Me	CH <sub>2</sub> OMe
38-3	38-4	Me	CH <sub>2</sub> OEt
39-3	39-4	Me	O-i-Pr
40-3	40-4	Me	O-n-Bu
41-3	41-4	Me	OCH <sub>2</sub> CF <sub>3</sub>

Compound 3-3 was separated into its enantiomers R3-3 and S3-3

Table 6



Compounds of formula (XX) wherein R<sub>201</sub> is cyano; R<sub>202</sub> is S(O)<sub>2</sub>CF<sub>3</sub>; R<sub>204</sub> is N(R<sub>205</sub>)C(O)CR<sub>206</sub>R<sub>207</sub>R<sub>208</sub>; R<sub>207</sub> and R<sub>208</sub> = H; R<sub>211</sub> is Cl, X<sub>1</sub> is C-Cl, and R<sub>213</sub> is CF<sub>3</sub> or SF<sub>5</sub>.

Compound Number (R <sub>213</sub> = CF <sub>3</sub> )	Compound Number (R <sub>213</sub> = SF <sub>5</sub> )	R <sub>205</sub>	R <sub>206</sub>
1-5	1-6	Me	H
2-5	2-6	Me	OMe
3-5	3-6	Me	OEt
4-5	4-6	Me	O-i-Pr
5-5	5-6	Me	O-n-Bu
6-5	6-6	Et	H
7-5	7-6	Et	OMe
8-5	8-6	Et	OEt
9-5	9-6	Et	O-i-Pr
10-5	10-6	Et	O-n-Bu
11-5	11-6	n-Pr	H
12-5	12-6	n-Pr	OMe
13-5	13-6	n-Pr	OEt
14-5	14-6	n-Pr	O-i-Pr
15-5	15-6	n-Pr	O-n-Bu
16-5	16-6	i-Pr	H
17-5	17-6	i-Pr	OMe
18-5	18-6	i-Pr	OEt
19-5	19-6	i-Pr	O-i-Pr
20-5	20-6	i-Pr	O-n-Bu
21-5	21-6	n-Bu	H
22-5	22-6	n-Bu	OMe
23-5	23-6	n-Bu	OEt
24-5	24-6	n-Bu	O-i-Pr
25-5	25-6	n-Bu	O-n-Bu
26-5	26-6	CH <sub>2</sub> Cy	H
27-5	27-6	CH <sub>2</sub> Cy	OMe

28-5	28-6	CH <sub>2</sub> Cy	OEt
29-5	29-6	CH <sub>2</sub> Cy	O-i-Pr
30-5	30-6	CH <sub>2</sub> Cy	O-n-Bu
31-5	31-6	CH <sub>2</sub> CCH	H
32-5	32-6	CH <sub>2</sub> CCH	OMe
33-5	33-6	CH <sub>2</sub> CCH	OEt
34-5	34-6	CH <sub>2</sub> CCH	O-i-Pr
35-5	35-6	CH <sub>2</sub> CCH	O-n-Bu
36-5	36-6	Me	OAc
37-5	37-6	Me	CH <sub>2</sub> OMe
38-5	38-6	Me	CH <sub>2</sub> OEt
39-5	39-6	Me	O-i-Pr
40-5	40-6	Me	O-n-Bu
41-5	41-6	Me	OCH <sub>2</sub> CF <sub>3</sub>

Table 7

Compounds of formula (XX) wherein R<sub>201</sub> is cyano; R<sub>202</sub> is

SCF<sub>3</sub>; R<sub>204</sub> is N(R<sub>205</sub>)C(O)CR<sub>206</sub>R<sub>207</sub>R<sub>208</sub>; R<sub>211</sub> is Cl; X<sub>1</sub> is C-Cl;

5 and R<sub>213</sub> is CF<sub>3</sub> or SF<sub>5</sub>-

Compound Number (R <sub>213</sub> =CF <sub>3</sub> )	Compound Number (R <sub>213</sub> =SF <sub>5</sub> )	R <sub>205</sub>	R <sub>206</sub>	R <sub>207</sub> , R <sub>208</sub>
1-7	1-8	Me	H	-CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> O-
2-7	2-8	Et	H	-CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> O-
3-7	3-8	i-Pr	H	-CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> O-
4-7	4-8	Pr	H	-CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> O-
5-7	5-8	Bu	H	-CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> O-
6-7	6-8	Cy	H	-CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> O-
7-7	7-8	CH <sub>2</sub> Cy	H	-CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> O-

Compound 1-7 was also separated into its enantiomers, called (R)1-7 and (S)1-7.

Table 8

Compounds of formula (XX) wherein R<sub>201</sub> is cyano; R<sub>202</sub> is S(O)CF<sub>3</sub>; R<sub>204</sub> is N(R<sub>205</sub>)C(O)CR<sub>206</sub>R<sub>207</sub>R<sub>208</sub>; R<sub>211</sub> is Cl, X<sub>1</sub> is C-Cl; and R<sub>213</sub> is CF<sub>3</sub> or SF<sub>5</sub>.

Compound Number (R <sub>213</sub> =CF <sub>3</sub> )	Compound Number (R <sub>213</sub> =SF <sub>5</sub> )	R <sub>205</sub>	R <sub>206</sub>	R <sub>207</sub> , R <sub>208</sub>
1-9	1-10	Me	H	-CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> O-
2-9	2-10	Et	H	-CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> O-
3-9	3-10	i-Pr	H	-CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> O-
4-9	4-10	Pr	H	-CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> O-
5-9	5-10	Bu	H	-CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> O-
6-9	6-10	CH <sub>2</sub> Cy	H	-CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> O-
7-9	7-10	Cy	H	-CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> O-

- 5 Compound 1-9 was separated into its diastereomers, (R,R)1-9, (S,R)1-9, (S,S)1-9, (R,S)1-9. The first designation of absolute configuration refers to the configuration of the sulfoxide moiety, the second to the chiral carbon.

Table 9

- 10 Compounds of formula (XX) wherein R<sub>201</sub> is cyano; R<sub>202</sub> is S(O)<sub>2</sub>CF<sub>3</sub>; R<sub>204</sub> is N(R<sub>205</sub>)C(O)CR<sub>206</sub>R<sub>207</sub>R<sub>208</sub>; R<sub>211</sub> is Cl; X<sub>1</sub> is C-Cl; and R<sub>213</sub> is CF<sub>3</sub> or SF<sub>5</sub>.

Compound Number (R <sub>213</sub> =CF <sub>3</sub> )	Compound Number (R <sub>213</sub> =SF <sub>5</sub> )	R <sub>205</sub>	R <sub>206</sub>	R <sub>207</sub> , R <sub>208</sub>
1-11	1-12A	Me	H	-CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> O-
2-11	2-12A	Et	H	-CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> O-
3-11	3-12A	i-Pr	H	-CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> O-
4-11	4-12A	Pr	H	-CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> O-
5-11	5-12A	Bu	H	-CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> O-
6-11	6-12A	Cy	H	-CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> O-
7-11	7-12A	CH <sub>2</sub> Cy	H	-CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> O-

Compound 1-11 was also separated into its diastereomers, (R)1-11 and (S)1-11.

Table 10

Compounds of formula (XX) wherein R<sub>201</sub> is cyano; R<sub>204</sub> is  
 5 N(R<sub>205</sub>)C(O)CR<sub>206</sub>R<sub>207</sub>R<sub>208</sub>; R<sub>207</sub> and R<sub>208</sub> are H; R<sub>211</sub> is Cl, X<sub>1</sub>  
 is C-Cl; and R<sub>213</sub> is CF<sub>3</sub> or SF<sub>5</sub>.

Compound Number (R <sub>213</sub> =CF <sub>3</sub> )	Compound Number (R <sub>213</sub> =SF <sub>5</sub> )	R <sub>205</sub>	R <sub>206</sub>
R <sub>202</sub> =SCF <sub>3</sub>			
1-12	1-13	Cy	H
2-12	2-13	Cy	OMe
3-12	3-13	Cy	OEt
4-12	4-13	Cy	i-O-Pr
5-12	5-13	Cy	O-n-Bu
R <sub>202</sub> =S(O)CF <sub>3</sub>			
6-12	6-13	Cy	H
7-12	7-13	Cy	OMe
8-12	8-13	Cy	OEt
9-12	9-13	Cy	O-i-Pr
10-12	10-13	Cy	O-n-Bu
R <sub>202</sub> =S(O) <sub>2</sub> CF <sub>3</sub>			
11-12	11-13	Cy	H
12-12	12-13	Cy	OMe
13-12	13-13	Cy	OEt
14-12	14-13	Cy	O-i-Pr
15-12	15-13	Cy	O-n-Bu

10

Table 11

Compounds of formula (XX) wherein R<sub>201</sub> is cyano; R<sub>204</sub> is

-N(R<sub>205</sub>)C(O)OR<sub>207</sub>; R<sub>211</sub> is Cl; X<sub>1</sub> is C-Cl, and R<sub>213</sub> is CF<sub>3</sub>

or SF<sub>5</sub>.

Compound Number (R <sub>213</sub> =CF <sub>3</sub> )	Compound Number (R <sub>213</sub> =SF <sub>5</sub> )	R <sub>205</sub>	R <sub>207</sub>
R <sub>202</sub> is SCF <sub>3</sub>			
67-1	67-2	Me	Me
68-1	68-2	Me	Et
69-1	69-2	Me	i-Pr
70-1	70-2	Me	n-Pr
71-1	71-2	Et	Me
72-1	72-2	Et	Et
73-1	73-2	Et	i-Pr
74-1	74-2	Et	n-Pr
75-1	75-2	n-Pr	Me
76-1	76-2	n-Pr	Et
77-1	77-2	n-Pr	i-Pr
78-1	78-2	n-Pr	n-Pr
79-1	79-2	i-Pr	Me
80-1	80-2	i-Pr	Et
81-1	81-2	i-Pr	i-Pr
82-1	82-2	i-Pr	n-Pr
R <sub>202</sub> is S(O)CF <sub>3</sub>			
83-1	83-2	Me	Me
84-1	84-2	Me	Et
85-1	85-2	Me	i-Pr
86-1	86-2	Me	n-Pr
87-1	87-2	Et	Me
88-1	88-2	Et	Et
89-1	89-2	Et	i-Pr
90-1	90-2	Et	n-Pr
91-1	91-2	n-Pr	Me
92-1	92-2	n-Pr	Et
93-1	93-2	n-Pr	i-Pr

94-1	94-2	n-Pr	n-Pr
95-1	95-2	i-Pr	Me
96-1	96-2	i-Pr	Et
97-1	97-2	i-Pr	i-Pr
98-1	98-2	i-Pr	n-Pr
R <sub>202</sub> is S(O) <sub>2</sub> CF <sub>3</sub>			
99-1	99-2	Me	Me
100-1	100-2	Me	Et
101-1	101-2	Me	i-Pr
102-1	102-2	Me	n-Pr
103-1	103-2	Et	Me
104-1	104-2	Et	Et
105-1	105-2	Et	i-Pr
106-1	106-2	Et	n-Pr
107-1	107-2	n-Pr	Me
108-1	108-2	n-Pr	Et
109-1	109-2	n-Pr	i-Pr
110-1	110-2	n-Pr	n-Pr
111-1	111-2	i-Pr	Me
112-1	112-2	i-Pr	Et
113-1	113-2	i-Pr	i-Pr
114-1	114-2	i-Pr	n-Pr

Table 12

Compounds of formula (XX) wherein R<sub>201</sub> is cyano; R<sub>202</sub> is S(O)<sub>h</sub>CF<sub>3</sub>; R<sub>204</sub> is N(R<sub>205</sub>)C(O)CR<sub>206</sub>R<sub>207</sub>R<sub>208</sub>; R<sub>211</sub> is Cl; X<sub>1</sub> is C-Cl, and R<sub>213</sub> is CF<sub>3</sub> or SF<sub>5</sub>.

Compound Number (R <sub>213</sub> = CF <sub>3</sub> )	Compound Number (R <sub>213</sub> = SF <sub>5</sub> )	h	R <sub>205</sub>	R <sub>206</sub>	R <sub>207</sub>	R <sub>208</sub>
115-1	115-2	0	Me	H	Me	Me
116-1	116-2	0	Me	OEt	H	Me
117-1	117-2	0	Me	H	cyclopropyl	
118-1	118-2	0	Me	OMe	H	Me

119-1	119-2	0	Me	OEt	Me	Me
120-1	120-2	2	Me	OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H
121-1	121-2	0	Me	H	-CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> O-	
122-1	122-2	1	Me	OEt	H	Me
123-1	123-2	0	Me	H	H	Me
124-1	124-2	0	Me	H	H	Et

Table 13

Compounds of formula (XX) wherein R<sub>201</sub> is cyano; R<sub>202</sub> is S(O)<sub>h</sub>CF<sub>3</sub>; R<sub>204</sub> is N(R<sub>205</sub>)C(O)aryl; R<sub>211</sub> is Cl; X<sub>1</sub> is C-Cl, R<sub>205</sub> is CH<sub>3</sub>; and R<sub>213</sub> is CF<sub>3</sub> or SF<sub>5</sub>. Within this table the following symbols are defined:

Ph means phenyl; Fu means furyl

Th means the thiophene radical

Pyr means pyridyl

Compound Number (R <sub>213</sub> =CF <sub>3</sub> )	Compound Number (R <sub>213</sub> =SF <sub>5</sub> )	aryl
R <sub>202</sub> is SCF <sub>3</sub>		
125-1	125-2	Ph
126-1	126-2	4-OMe-Ph
127-1	127-2	4-CF <sub>3</sub> -Ph
128-1	128-2	2-Th
129-1	129-2	3-Th
130-1	130-2	2-Fu
131-1	131-2	3-Fu
132-1	132-2	2-Pyr
133-1	133-2	3-Pyr
134-1	134-2	4-Pyr
135-1	135-2	6-Cl-2-Pyr
136-1	136-2	6-CF <sub>3</sub> -2-Pyr
137-1	137-2	5-Cl-2-Fu
138-1	138-2	5-CF <sub>3</sub> -2-Fu
139-1	139-2	5-OMe-2-Th
140-1	140-2	5-CF <sub>3</sub> -2-Th

R <sub>202</sub> is S(O)CF <sub>3</sub>		
125-3	125-4	Ph
126-3	126-4	4-OMe-Ph
127-3	127-4	4-CF <sub>3</sub> -Ph
128-3	128-4	2-Th
129-3	129-4	3-Th
130-3	130-4	2-Fu
131-3	131-4	3-Fu
132-3	132-4	2-Pyr
133-3	133-4	3-Pyr
134-3	134-4	4-Pyr
135-3	135-4	6-Cl-2-Pyr
136-3	136-4	6-CF <sub>3</sub> -2-Pyr
137-3	137-4	5-Cl-2-Fu
138-3	138-4	5-CF <sub>3</sub> -2-Fu
139-3	139-4	5-OMe-2-Th
140-3	140-4	5-CF <sub>3</sub> -2-Th
R <sub>202</sub> is S(O) <sub>2</sub> CF <sub>3</sub>		
125-5	125-6	Ph
126-5	126-6	4-OMe-Ph
127-5	127-6	4-CF <sub>3</sub> -Ph
128-5	128-6	2-Th
129-5	129-6	3-Th
130-5	130-6	2-Fu
131-5	131-6	3-Fu
132-5	132-6	2-Pyr
133-5	133-6	3-Pyr
134-5	134-6	4-Pyr
135-5	135-6	6-Cl-2-Pyr
136-5	136-6	6-CF <sub>3</sub> -2-Pyr
137-5	137-6	5-Cl-2-Fu
138-5	138-6	5-CF <sub>3</sub> -2-Fu
139-5	139-6	5-OMe-2-Th
140-5	140-6	5-CF <sub>3</sub> -2-Th



The present invention also relates to a composition comprising a parasitically effective, substantially non-emetic amount of a compound of formula (I) or a salt thereof or a compound of formula (XX) or a salt thereof and an acceptable carrier. Acceptable carriers for the use of the compounds are generally known to the skilled addressee concerned with pest control in animals, particularly domestic animals, most preferably dogs or cats.

The compositions which can be used in the invention can comprise generally from about 0.001 to 95% of the compound of formula (I) or a salt thereof or a compound of formula (XX) or a salt thereof. The remainder of the composition up to 100% comprises a carrier as well as generally various additives. In this specification and the accompanying claims, percentages are by weight.

The diluted liquid formulations generally comprise from about 0.001 to about 3% of compound of formula (I) or a salt thereof or a compound of formula (XX) or a salt thereof, preferably from about 0.1 to about 0.5%.

Solid formulations generally comprise from about 0.1 to about 8% of compound of formula (I) or a salt thereof or a compound of formula (XX) or a salt thereof, preferably from about 0.5 to about 1.5%.

Compositions for oral administration comprise one or more of the compounds of general formula (I) or salts thereof or compounds of formula (XX) or salts thereof in association with veterinarily acceptable carriers or coatings and include, for example, tablets, pills, capsules, gels, drenches, medicated feeds, medicated drinking water, medicated dietary supplements, slow-release boluses or other slow-release devices intended to be retained within the gastro-intestinal tract. Any of these may incorporate the active ingredients contained within micro-capsules or coated with acid-labile or alkali-labile or other pharmaceutically acceptable enteric coatings. Feed premixes or concentrates containing

compounds of the present invention for use in preparation of medicated diets, drinking water or other materials for consumption by animals may also be used. In a highly preferred embodiment, the compositions are administered postprandially, preferably from just after a meal to 2 hours after the meal.

In a highly preferred embodiment, there is provided a product which is readily chewed by the animal and which product does generally not allow human contamination when the product is provided to the animal by hand.

The compounds of general formula (I) or salts thereof or compounds of formula (XX) or salts thereof may be administered before, during or after meals. The compounds of general formula (I) or salts thereof or compounds of formula (XX) or salts thereof may be mixed with a carrier and/or a foodstuff.

According to the present invention the compound of formula (I) or a salt thereof or a compound of formula (XX) or a salt thereof is administered orally in a dose to the animal in a dose range generally from 0.1 to 500 mg/kg of the compound of formula (I) or a salt thereof or a compound of formula (XX) or a salt thereof per kilogram of animal body weight (mg/kg), preferably from 1 to 100 mg/kg, more preferably from 1 to 50 mg/kg, even more preferably from 2 to 25 mg/kg, most preferably from 3 to 15 mg/kg.

According to the present invention, the frequency of treatment of the animal, preferably the domestic animal to be treated by the compound of formula (I) or a salt thereof or a compound of formula (XX) or a salt thereof is generally from about once per week to about once per year, preferably from about once every two weeks to about once every six months, more preferably from about once every two weeks to once every three months, and most preferably from about once every two weeks to about once every six weeks.

Generally the animal to be treated is a domestic animal, preferably a domestic companion animal. More preferably the animal to be treated is a dog and/or a cat.

5 The compounds of the invention may be administered most advantageously with another parasitically effective material, such as an endoparasiticide, and/or an ectoparasiticide, and/or an endectoparasiticide. For example, such compounds include macrocyclic lactones such as avermectins or milbemycins e.g., ivermectin; pyratel (generally administered as pyrantel pamoate) or an insect growth  
10 regulator such as lufenuron or methoprene.

By the term "parasites" as used in the specification and claims is meant endoparasites and ectoparasites of warm-blooded animals, particularly ectoparasites. Preferably, fleas and/or ticks are controlled by the method of the present invention.

15 Illustrative of specific parasites of various host animals which may be controlled by the method of this invention include arthropods such as:

Mites: Mesostigmata spp. e.g. mesostigmatids such as the chicken mite, Dermanyssus gallinae; itch or scab mites such as Sarcoptidae spp. for example Sarcoptes scabiei; mange mites such as Psoroptidae spp. including Chorioptes bovis and Psoroptes ovis; chiggers e.g. Trombiculidae spp. for example the north american chigger, Trombicula alfreddugesi;

25 Ticks: e.g., soft-bodied ticks including Argasidae spp. for example Argas spp. and Ornithodoros spp ; hard-bodied ticks including Ixodidae spp., for example Rhipicephalus sanguineus, and Boophilus spp.;

Lice: sucking lice, e.g., Menopon spp. and Bovicola spp.; biting lice, e.g., Haematopinus spp., Linognathus spp. and Solenopotes spp.;

30 Fleas: e.g., Ctenocephalides spp., such as dog flea (Ctenocephalides canis) and cat flea (Ctenocephalides felis); Xenopsylla

spp. such as oriental rat flea [Xenopsylla cheopis]; and Pulex spp. such as human flea [Pulex irritans];

5 True bugs: e.g., Cimicidae or including the common bed bug (Cimex lectularius);, Triatominae spp. including triatomid bugs also known as kissing bugs; for example Rhodnius prolixus and Triatoma spp.;

bloodsucking adult flies: (e.g., horn fly [Haematobia irritans], horse fly [Tabanus spp.], stable fly [Stomoxys calcitrans], black fly [Simulium spp.], deer fly [Chrysops spp.], louse fly [Melophagus 10 ovinus], tsetse fly [Glossina spp.], mosquitoes [Culex spp., Anopheles spp., and Aedes spp.); and

parasitic fly maggots: (e.g., bot fly [Oestrus ovis and Cuterebra spp.], blow fly [Phaenicia spp.], screwworm [Cochliomyia 15 hominivorax], cattle grub [Hypoderma spp.], fleeceworm.

The present invention also relates to a use of a compound of formula (I) or a salt thereof or a compound of formula (XX) or a salt thereof hereinbefore described as a therapeutic agent, preferably for animals, more preferably for domestic animals.

20

The veterinary composition may be sterile or non-sterile. It may be a liquid (e.g. aqueous) or solid (e.g., dry) composition, in particular a freeze-dried composition, which, by addition of water or another liquid, orally effective solutions may be prepared.

25

The present invention also relates to a use of a compound of formula (I) or a salt thereof or a compound of formula (XX) or a salt thereof as hereinbefore defined for the manufacture of a veterinary composition for the control of parasites in or on an animal.

The present invention also relates to a method of cleaning animals in good health comprising the application to the animal of a

compound of formula (I) or a salt thereof or a compound of formula (XX) or a salt thereof as hereinbefore defined to the animal.

The method of cleaning an animal is not a method of treatment by therapy of the animal body per se, because


5 (a) the animal is in good health and requires no substantial treatment to correct a deficiency of health;

(b) the cleaning of the animal is not intended to be done by veterinary personnel, but by persons interested in the cleaning of the animal; and

10 (c) the purpose of such cleaning is to avoid unpleasant conditions for humans and the environment in which humans inhabit so as to not infest the said humans with arthropods carried by the animal.

By "carrier" is meant an organic or inorganic material, which can be natural or synthetic, and which is associated with the compound and which facilitates its application to the animal. This carrier is thus  
15 generally inert and should be arthropodically acceptable. The carrier can be solid (e.g., clay, silicates, silica, resins, wax.) or liquid (e.g., water, alcohols, ketones, oil solvents, polar aprotic solvents) An example of an oil solvent is corn oil. An example of a polar aprotic  
20 solvent is dimethyl sulfoxide.


INCORPORERThe compounds of formula (II) wherein R<sub>21</sub>, R<sub>22</sub>, R<sub>24</sub>, R<sub>31</sub>, R<sub>33</sub> and X are as defined above may be prepared from the compounds of formula (III):



(III)

wherein  $R_{21}$ ,  $R_{22}$ ,  $R_{31}$ ,  $R_{33}$  and  $X$  are as defined above, using processes described in European Patent Publications 0511845 or 0659745.

5 According to a feature of the present invention, compounds of formula (II) wherein  $R_{21}$ ,  $R_{22}$ ,  $R_{31}$ ,  $R_{33}$  and  $X$  are as defined above and  $R_{24}$  is  $-N=C(R_{25})-NR_{26}R_{27}$  wherein  $R_{25}$ ,  $R_{26}$  and  $R_{27}$  are as defined above may be prepared by reacting a compound of formula (III) with a compound of formula (IV):




(IV)

10 wherein  $R_{25}$ ,  $R_{26}$  and  $R_{27}$  are as defined above and  $R_{100}$  is generally an alkyl group. The reaction is optionally conducted in the presence of a catalyst such as a mineral or organic acid (for example hydrochloric acid), generally using from 1 to 100 equivalents of (IV), preferably using 1 to 10 equivalents of (IV), and is preferably conducted in an organic solvent such as tetrahydrofuran, toluene, or N,N-dimethylformamide, at a temperature of from 0°C to 150°C. Additional  
 15  
 20 adjuvants such as drying agents (e.g. magnesium sulfate, potassium carbonate, or molecular sieves) may also be advantageous to the reaction. Compounds of formula (IV) are known or may be prepared by known procedures.


25 According to a feature of the present invention, compounds of formula (II) wherein  $R_{21}$ ,  $R_{22}$ ,  $R_{31}$ ,  $R_{33}$  and  $X$  are as defined above and  $R_{24}$  is  $-N=C(R_{25})-NR_{26}R_{27}$  wherein  $R_{25}$ ,  $R_{26}$  and  $R_{27}$  are as

defined above, may be prepared by the reaction of a compound of formula (V):



(V)


wherein R<sub>21</sub>, R<sub>22</sub>, R<sub>25</sub>, R<sub>31</sub>, R<sub>33</sub>, X and R<sub>100</sub> are as defined  
 5 above, with a compound of formula (VI):



(VI)

wherein R<sub>26</sub> and R<sub>27</sub> are as defined above. The reaction is  
 generally conducted using the same conditions as used for the  
 preparation of compounds of formula (II) by the reaction of compounds  
 10 of formula (III) with compounds of formula (IV).

According to a feature of the present invention, compounds of  
 formula (II) wherein R<sub>24</sub> is -N=C(R<sub>25</sub>)-NR<sub>27</sub>R<sub>28</sub>, and R<sub>21</sub>, R<sub>22</sub>, R<sub>25</sub>,  
 R<sub>27</sub>, R<sub>31</sub>, R<sub>33</sub> and X are as defined above, and R<sub>28</sub> is COR<sub>34</sub> wherein  
 R<sub>34</sub> is as defined above, may be prepared by the reaction of the  
 15 corresponding compounds of formula (II) wherein R<sub>24</sub> is -N=C(R<sub>25</sub>)-  
 NR<sub>27</sub>H with an acid chloride of formula (VII):



wherein R<sub>34</sub> is as defined above. The reaction is generally  
 performed in the presence of a base such as a trialkylamine for example

triethylamine in a solvent such as dichloromethane, at a temperature of from 0°C to 50°C.

5

According to a feature of the present invention, compounds of formula (II) wherein  $R_{24}$  is  $-N=C(R_{25})-NR_{27}R_{28}$ , and  $R_{21}$ ,  $R_{22}$ ,  $R_{25}$ ,  $R_{27}$ ,  $R_{31}$ ,  $R_{33}$  and  $X$  are as defined above and  $R_{28}$  is  $-S(O)_tR_{30}$  may be prepared by the reaction of the corresponding compound of formula (II) wherein  $R_{24}$  is  $-N=C(R_{25})-NR_{27}H$  with a sulfonyl chloride or a sulfenyl chloride of formula (VIII):

10



15

The reaction is generally performed in the presence of a weak base such as a trialkylamine for example triethylamine, or pyridine in a solvent such as dichloromethane, at a temperature of from 0°C to 50°C.

Compounds of formula (VI), (VII) and (VIII) are known or may be prepared by known procedures.

20


Compounds of formula (III) and (V) may be generally prepared according to known processes, for example as described in International Patent Publications WO 87/3781, WO 93/6089, WO 94/21606 WO 97/07102, WO 98/24767, , WO 98/28277, WO 98/28278 and WO 98/28279,, European Patent Application 295117, 846686, and United States Patent 5232940..

25

In another aspect of the present invention, compounds of formula (XX) wherein  $R_{204}$  is  $-N(R_{205})C(O)CR_{206}R_{207}R_{208}$ ,  $N(R_{205})C(O)aryl$ , or  $N(R_{205})C(O)OR_{207}$  are generally prepared from compounds of formula (XXI):

30






(XXI)

respectively by reaction with halides of formulae

$X_2C(O)CR_{206}R_{207}R_{208}$ ,  $X_2C(O)aryl$ , or  $X_2C(O)OR_{207}$ , wherein

$R_{201}$ ,  $R_{202}$ ,  $R_{205}$ ,  $R_{206}$ ,  $R_{207}$ ,  $R_{208}$ ,  $R_{211}$ ,  $R_{213}$ , and  $X_1$  are defined above and wherein  $X_2$  is a halogen atom. The reaction is generally carried out in the presence of a base, generally using from 1 to 10 molar equivalents of the halide, and is preferably conducted in the presence of an organic solvent such as tetrahydrofuran, methylene chloride, at a temperature of from  $0^\circ\text{C}$  to  $150^\circ\text{C}$ .

Compounds of formula (XXI) may be prepared from a compound of formula (XXII):




(XXII)

by reaction with a compound of formula (XXIII):

$X_2R_{205}(XXIII)$

wherein  $R_{201}$ ,  $R_{202}$ ,  $R_{205}$ ,  $R_{211}$ ,  $R_{213}$ ,  $X_1$  and  $X_2$  are defined above. Compounds of formula (XXIII) are generally known in the art as alkylhalides or substituted alkylhalides. Compounds of formula XXII may be prepared by methods described in International Patent Publications WO 87/3781, WO 93/6089, WO 94/21606, WO 97/07102, WO 98/24767, , WO 98/28277, WO 98/28278 and WO 98/28279, European Patent Application 295117, 659745, 846686, and United States Patent 5232940 or other methods known to the person skilled in the art.

Alternatively compounds of formula (XXI) may be prepared by reduction of compounds of formula (XXIV):



(XXIV)

wherein  $R_{201}$ ,  $R_{202}$ ,  $R_{211}$ ,  $R_{213}$  and  $X_1$  are defined above. The reduction generally is effected by the use of a standard hydride ion donor, for example sodium borohydride or sodium cyanoborohydride. The reaction is generally effected in a polar solvent such as ethanol or

methanol and generally using from 1 to 10 molar equivalents of the hydride, and is preferably conducted at temperature of from -100°C to 150°C.

5       Compounds of formula (XXIV) may be prepared using methods described in EP 295117, WO 97/22593 or other methods known to those skilled in the art.

10       In another aspect of the invention there are provided the compounds 3-Cyano-1-(2,6-dichloro-4-trifluoromethylphenyl)-5-ethoxymethylideneamino-4-trifluoromethylsulfinylpyrazole and 3-cyano-1-(2,6-dichloro-4-trifluoromethylphenyl)-5-methylamino-4-trifluoromethylsulfinylpyrazole which are useful intermediates for the preparation of compounds for use according to the present invention.

#### Biological Example

20

Compounds 1-1, 2-1, 3-1, 4-1, 11-1, 13-1, 28-1, 31-1, 32-1, 36-1, 37-1, 38-1, 1-3, 2-3, 3-3, 4-3, 6-3, 41-3, 1-5, 2-5, 3-5, 6-5, 11-5, 27-5, 28-5, 1-7, 3-7, 5-7, 1-9, 1-11, 6-11, 7-11, 1-12, 11-12, 13-12, 67-1, 68-1, 69-1, 70-1, 72-1, 75-1, 76-1, 77-1, 78-1, 79-1, 80-1, 81-1, 82-1, 115-1, 25   116-1, 117-1, 118-1, 119-1, 120-1, 121-1, 122-1, 123-1, 124-1, 211-1, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, R3-3, S3-3, R1-7, S1-7, (R,S)1-9, (R,R)1-9, (S,R)1-9, (S,S) 1-9, S1-11, R1-11, 126-1, 127-1 and 130-1 were formulated as a 30 mg/mL formulations in a 1:1 volume/volume solution of dimethyl sulfoxide and corn oil. Using this formulation, mixed breed dogs and cats were 30   treated at a rate of 10 mg of the compound per kg (mg/kg) of body weight of the dog and 20 mg/kg of the cat treated. The animals were fasted for at least 8 hours prior to treatment, fed half of the daily ration

immediately prior to treatment, then allowed access to the remainder of the daily ration immediately following treatment.

5 All dogs were infested with cat fleas (*Ctenocephalides felis*) and with ticks (*Rhipicephalus sanguineus*) 1 day prior to administration of the compound. Cats were only infested with fleas. The initial flea and tick counts were performed 1 day after the administration of the compounds. At 7, 14, 21 and 28 days after treatment the dogs were re-infested with ticks and 8, 15, 22 and 29 days after treatment the dogs and cats were re-infested with fleas. At 1, 9, 16, 23 and 30 days after  
10 treatment the control of fleas and ticks in treated dogs and cats was determined versus a group of infested dogs and cats which received a placebo consisting of a 1:1 volume/volume solution of dimethyl sulfoxide and corn oil. To determine the efficacies of the compounds, the arthropods were combed from the animals and counted.

15 In the animals treated with the compounds above, there was substantially no emesis after 2, 8 and 24 hours. Generally long-term control of fleas and ticks was provided in dogs. In the cats treated, there was commercially acceptable control of fleas for at least one day post treatment.

20 The results of this example were superior to those obtained with compounds of the prior art, for example, fipronil.

The following non-limiting Synthesis Examples illustrate the preparation of compounds of formula (I) and the Reference Examples  
25 illustrate the preparation of intermediates used in their synthesis.

#### Synthesis Example 1

A solution of 5-amino-3-cyano-1-(2,6-dichloro-4-trifluoromethylphenyl)-4-trifluoromethylthiopyrazole (1 g) in N,N-dimethylformamide dimethyl acetal was heated at 50°C for 1 hour.  
30

Evaporation of solvents gave 3-cyano-1-(2,6-dichloro-4-trifluoromethylphenyl)-5-N'-dimethylaminomethylideneamino-4-trifluoromethylthiopyrazole m.p.141°C.

By proceeding in a similar manner the following compounds were also prepared:

3-cyano-1-(2,6-dichloro-4-trifluoromethylphenyl)-5-N'-dimethylaminomethylideneamino-4-trifluoromethylsulfonylpyrazole m.p.209°C; and

3-cyano-1-(2,6-dichloro-4-trifluoromethylphenyl)-5-N'-dimethylaminomethylideneamino-4-trifluoromethylsulfinylpyrazole m.p.207°C.

### Synthesis Example 2

A solution of 3-cyano-1-(2,6-dichloro-4-trifluoromethylphenyl)-5-ethoxymethylideneamino-4-trifluoromethylthiopyrazole (5 g) in ethanol was treated with benzylamine ( 11.4 ml), stirred overnight, evaporated and purified by reverse-phase column chromatography (C-18 stationary phase column, eluting with MeOH/water) to give 5-N'-benzylaminomethylideneamino-3-cyano-1-(2,6-dichloro-4-trifluoromethylphenyl)- 4-trifluoromethylthiopyrazole 1.18 g), m.p.113°C.

By proceeding in a similar manner the compounds of formula (II) wherein R<sub>21</sub> is CN; R<sub>24</sub> is -N=CH-NHR<sub>26</sub>; R<sub>31</sub> is Cl; X is C-Cl; and R<sub>33</sub> is CF<sub>3</sub> shown in Table 4 were also prepared.

**Table 4**

Compd No.	R22	R26	M.P. °C
21	SCF3	CH2CN	175
211-1	SCF3	CH2CF3	130
222	SCF3	CH3	173
223	SOCF3	CH2Ph	173
225	SOCF3	CH3	144
226	SOCF3	CH3	144

227	SOCF <sub>3</sub>	CH <sub>2</sub> CF <sub>3</sub>	175
228	SO <sub>2</sub> CF <sub>3</sub>	2-propynyl	149
229	SO <sub>2</sub> CF <sub>3</sub>	CH <sub>2</sub> Ph	182
230	SO <sub>2</sub> CF <sub>3</sub>	CH <sub>2</sub> CF <sub>3</sub>	183
209-1	SCF <sub>3</sub>	iPr	
207-1	SCF <sub>3</sub>	CH <sub>2</sub> CH <sub>3</sub>	141

### Reference Example 1

A solution of 5-amino-3-cyano-1-(2,6-dichloro-4-  
 trifluoromethylphenyl)-4-trifluoromethylthiopyrazole (500 g) in triethyl  
 orthoformate was treated with concentrated hydrochloric acid (10 ml)  
 and heated at 50°C. After 8 hours the reaction mixture was evaporated  
 to give a solid which was washed (heptane) and air-dried to give 3-  
 cyano-1-(2,6-dichloro-4-trifluoromethylphenyl)-5-  
 ethoxymethylideneamino-4-trifluoromethylthiopyrazole (217 g), m.p.  
 68°C.

By proceeding in a similar manner the following intermediates were also prepared:

3-cyano-1-(2,6-dichloro-4-trifluoromethylphenyl)-5-  
 ethoxymethylideneamino-4-trifluoromethylsulfinylpyrazole, m.p. 63°C;  
 and 3-cyano-1-(2,6-dichloro-4-trifluoromethylphenyl)-5-  
 ethoxymethylideneamino-4-trifluoromethylsulfonylpyrazole,  
 m.p. 118°C.

### Synthesis Example 3

3-Cyano-1-(2,6-dichloro-4-trifluoromethylphenyl)-5-  
 methylamino-4-trifluoromethylsulfinylpyrazole (111.55 g, 0.247 moles),  
 triethylamine (62.45 g, 0.618 moles), 4-dimethylaminopyridine (3 g,  
 0.0247 moles), and tetrahydrofuran (700 ml) was combined. The  
 resulting solution was heated to 45°C and ethoxyacetyl chloride (45.2 g,

0.37 mmol) was added dropwise over 10 min. After 1 h, the mixture was evaporated to a brown residue, which was dissolved in 500 ml of ethyl acetate and washed with 2 x 300 ml of water. The organic phase was dried over magnesium sulfate, filtered, and evaporated to a brown oil. The oil was triturated with 1 L of hot cyclohexane. The resulting solids were collected by filtration and washed with 500 ml of hot cyclohexane, then air dried to afford of compound 3-3 as a beige powder (116.7 g). Evaporation of the mother liquors afforded a second crop of compound 3-3 (8.4 g).

In a similar fashion or by modifications according to methods known to the skilled addressee, the following compounds were prepared. The compound numbers in the left column refer to the Tables cited above.

Compound Number	Mass Spectral molecular ion + 1 (M+1)
1-1	477
2-1	507
3-1	521
4-1	535
11-1	505
13-1	549
28-1	537
31-1	517
32-1	531
36-1	535
37-1	521
38-1	535
1-3	593
2-3	523

3-3	537
4-3	551
6-3	491
41-3	473
1-5	509
2-5	539
3-5	553
6-5	523
11-5	537
27-5	579
28-5	593
1-7	533
3-7	561
5-7	575
1-9	549
1-11	565
6-11	591
7-11	605
1-12	535
11-12	565
13-12	579
67-1	493
68-1	507
69-1	521
70-1	521
72-1	521
75-1	521
76-1	535
77-1	549



78-1	549
79-1	521
80-1	535
81-1	549
82-1	549
115-1	505
116-1	535
117-1	503
118-1	521
119-1	549
120-1	583
121-1	547
122-1	551
123-1	491
124-1	505
126-1	507
127-1	607
130-1	529

**Reference Example 2:**

5

Step A: Preparation of 3-Cyano-1-(2,6-dichloro-4-trifluoromethylphenyl)-5-ethoxymethylideneamino-4-trifluoromethylsulfinylpyrazole.

10

A 12-L three-necked flask fitted with an overhead stirrer, heating mantle, water separator (e.g. Dean Stark trap) with condenser was placed under a nitrogen atmosphere and charged with 1.475 kg (3.37 moles) of fipronil and 6 L of triethyl orthoformate. The suspension was


heated to reflux over 2.5 h, then at reflux for 3 h with collection and removal of the distillate. The mixture was cooled to room temperature, then evaporated under reduced pressure at a bath temperature of 60-80°C, then at 50°C overnight. The resulting beige solid, 1.717 kg (95.8  
5 % by HPLC, 3.335 moles, 99% purity corrected yield) was used without further purification. (m.p. about 63°C)

Step B: Preparation of 3-Cyano-1-(2,6-dichloro-4-trifluoromethylphenyl)-5-methylamino-4-trifluoromethylsulfanylpyrazole.  
10

A 50 L reactor was charged with 3-Cyano-1-(2,6-dichloro-4-trifluoromethylphenyl)-5-ethoxymethylideneamino-4-trifluoromethylsulfanylpyrazole (1.645 g, 3.335 moles) and absolute  
15 ethanol (16 L) under nitrogen. The solution was cooled to 10°C, and sodium borohydride (266 g, 7.03 moles) was added slowly such that the temperature, remained generally below 35°C. After 6.75 h, some additional sodium borohydride (25 g, 0.66 moles) was added and stirring was continued overnight. Acetic acid (1.3 L, 22.7 moles) was  
20 added to quench, followed by 16 L of water. The resulting precipitate was collected by filtration, washed with water, and air-dried. Recrystallization from methanol afforded 3-Cyano-1-(2,6-dichloro-4-trifluoromethylphenyl)-5-methylamino-4-trifluoromethylsulfanylpyrazole (350 g) as an off-white solid. (m.p.  
25 about 227°C)

CLAIMS

- 5 1. A method of controlling parasites in or on an animal comprising administering orally to the animal a parasitically effective, substantially non-emetic amount of a 1-arylpyrazole of formula (I):



(I)

wherein:

- 10 R<sub>1</sub> is cyano, acetyl, C(S)NH<sub>2</sub>, alkyl, haloalkyl, C(=NOH)NH<sub>2</sub> or C(=NNH<sub>2</sub>)NH<sub>2</sub>;

R<sub>2</sub> is S(O)<sub>n</sub>R<sub>3</sub>; C<sub>2</sub>-C<sub>3</sub> alkenyl, C<sub>2</sub>-C<sub>3</sub> haloalkenyl, cycloalkyl, halocycloalkyl or C<sub>2</sub>-C<sub>3</sub> alkynyl;

R<sub>3</sub> is alkyl or haloalkyl;

- 15 R<sub>4</sub> is -N=C(R<sub>5</sub>)-Z-R<sub>6</sub>, -N=C(R<sub>5</sub>)-N(R<sub>7</sub>)-R<sub>8</sub>, or -N(R<sub>9</sub>)-C(R<sub>5</sub>)=NR<sub>6</sub>;

R<sub>5</sub> is hydrogen; alkyl; or alkyl substituted by halogen, alkoxy, haloalkoxy or -S(O)<sub>m</sub>R<sub>15</sub>;

- 20 R<sub>6</sub> and R<sub>7</sub> each independently represent hydrogen, alkyl, C<sub>3</sub>-C<sub>5</sub> alkenyl or C<sub>3</sub>-C<sub>5</sub> alkynyl; or

alkyl substituted by one or more halogen, alkoxy, haloalkoxy, amino, alkylamino, dialkylamino, cyano or -S(O)<sub>m</sub>R<sub>15</sub>; or alkyl

substituted by phenyl or pyridyl each of which is optionally substituted with one or more groups selected from halogen, nitro and alkyl; or

R<sub>6</sub> and R<sub>7</sub> may form together with the nitrogen to which they are attached a 3 to 7 membered ring which may additionally contain one or more heteroatoms selected from oxygen, nitrogen or sulfur;

R<sub>8</sub> is alkoxy, haloalkoxy, amino, alkylamino, dialkylamino, R<sub>14</sub>CO- or -S(O)<sub>t</sub>R<sub>10</sub>;

R<sub>9</sub>, R<sub>10</sub> and R<sub>14</sub> are alkyl or haloalkyl;

R<sub>11</sub> and R<sub>12</sub> are independently selected from halogen, hydrogen, CN and NO<sub>2</sub>;

R<sub>13</sub> is selected from halogen, haloalkyl, haloalkoxy, -S(O)<sub>q</sub>CF<sub>3</sub>, and -SF<sub>5</sub>;

R<sub>15</sub> is alkyl or haloalkyl;


X is selected from nitrogen and C-R<sub>12</sub>;

Z is O, S(O)<sub>a</sub>; or NR<sub>7</sub>;

a, m, n and q are independently selected from 0, 1, and 2; and

t is 0 or 2; and veterinarily acceptable salts thereof.

2. A method of controlling parasites in or on an animal comprising administering orally to the animal a parasitically effective, substantially non-emetic amount of a 1-arylpyrazole of formula (XX):



(XX)

wherein:

R<sub>201</sub> is cyano, C(O)alkyl, C(S)NH<sub>2</sub>, alkyl, C(=NOH)NH<sub>2</sub> or C(=NNH<sub>2</sub>)NH<sub>2</sub>;

5 R<sub>202</sub> is S(O)<sub>h</sub>R<sub>203</sub>, C<sub>2</sub>-C<sub>3</sub> alkenyl, C<sub>2</sub>-C<sub>3</sub> haloalkenyl, cycloalkyl, halocycloalkyl or C<sub>2</sub>-C<sub>3</sub> alkynyl;

R<sub>203</sub> is alkyl or haloalkyl;

R<sub>204</sub> is -N(R<sub>205</sub>)C(O)CR<sub>206</sub>R<sub>207</sub>R<sub>208</sub>, -N(R<sub>205</sub>)C(O)aryl, or -N(R<sub>205</sub>)C(O)OR<sub>207</sub>;

10 R<sub>205</sub> is alkyl, haloalkyl, cycloalkyl, halocycloalkyl, cycloalkylalkyl, halocycloalkylalkyl, alkoxyalkyl, haloalkoxyalkyl, C<sub>3</sub>-C<sub>3</sub> alkenyl, C<sub>3</sub>-C<sub>3</sub> haloalkenyl, C<sub>3</sub>-C<sub>3</sub> alkynyl, C<sub>3</sub>-C<sub>3</sub> haloalkynyl;

15 R<sub>206</sub> is hydrogen, halogen, alkoxy, haloalkoxy, alkoxyalkyl, haloalkoxyalkyl, formyloxy, alkylcarbonyloxy, haloalkylcarbonyloxy, alkylthio, haloalkylthio, alkylsulfinyl, haloalkylsulfinyl, alkylsulfonyl, haloalkylsulfonyl, alkylamino, dialkylamino, haloalkylamino, di(haloalkyl)amino, cycloalkyloxy, halocycloalkyloxy, alkoxyalkoxy, haloalkoxyalkoxy, alkoxyalkoxyalkoxy, aryloxy, or arylalkoxy;

20 R<sub>207</sub> and R<sub>208</sub> are independently hydrogen, alkyl, haloalkyl, cycloalkyl, or halocycloalkyl; or R<sub>207</sub> and R<sub>208</sub> may form together with the carbon to which they are attached a 3 to 7 membered ring which additionally may contain one or more heteroatoms selected from nitrogen, oxygen and sulfur;

X<sub>1</sub> is selected from nitrogen and C-R<sub>212</sub>;

25 R<sub>211</sub> and R<sub>212</sub> are independently selected from halogen, hydrogen, CN and NO<sub>2</sub>;

R<sub>213</sub> is selected from halogen, haloalkyl, haloalkoxy, -S(O)<sub>k</sub>CF<sub>3</sub>, and -SF<sub>5</sub>; and

h and k are independently selected from 0, 1, and 2;

and veterinarily acceptable salts thereof.

3. The method according to Claim 1 wherein the compound of formula (I) is that wherein:

5

$R_1$  is cyano or alkyl;

$R_2$  is  $S(O)_nR_3$ ;

$R_3$  is alkyl or haloalkyl;

$R_4$  is  $-N=C(R_5)-Z-R_6$ ;

$R_5$  is hydrogen, alkyl or haloalkyl;

10

$Z$  is O,  $S(O)_a$ ; or  $NR_7$ ;

$R_6$  and  $R_7$  are independently selected from hydrogen and unsubstituted or substituted alkyl; or

$R_6$  and  $R_7$  may form together with the nitrogen to which they are attached a 3 to 7 membered ring which may additionally contain one or more heteroatoms selected from oxygen, nitrogen or sulfur;  $X$  is selected from nitrogen and C- $R_{12}$ ;

15

$R_{11}$  and  $R_{12}$  are independently selected from halogen, hydrogen, CN and  $NO_2$ ;

20

$R_{13}$  is selected from halogen, haloalkyl, haloalkoxy,  $-S(O)_qCF_3$ , and  $-SF_5$ ;

$a$ ,  $n$  and  $q$  are independently selected from 0, 1, and 2.

4. The method according to claim 2 wherein the compound of formula (XX) is that wherein:

25

$R_{201}$  is cyano;

$R_{202}$  is  $S(O)_hR_{203}$ ;

$R_{203}$  is alkyl or haloalkyl;

$R_{204}$  is  $-N(R_{205})C(O)CR_{206}R_{207}R_{208}$ ;

R<sub>205</sub> is alkyl, haloalkyl, cycloalkyl, cycloalkylalkyl and  
halocycloalkylalkyl;

R<sub>206</sub> is alkoxy, haloalkoxy, or hydrogen;

R<sub>207</sub> and R<sub>208</sub> are independently hydrogen, alkyl, or haloalkyl; or

5 R<sub>207</sub> and R<sub>208</sub> may form together with the carbon to which they  
are attached a 3 to 7 membered ring which additionally may contain one  
or more heteroatoms selected from nitrogen, oxygen and sulfur;


X<sub>1</sub> is selected from nitrogen and C-R<sub>212</sub>;

10 R<sub>211</sub> and R<sub>212</sub> are independently selected from halogen,  
hydrogen, CN and NO<sub>2</sub>;

R<sub>213</sub> is selected from halogen, haloalkyl, haloalkoxy, -S(O)<sub>k</sub>CF<sub>3</sub>,  
and -SF<sub>3</sub>; and

h and k are independently selected from 0, 1, and 2.

15 5. A compound of formula (II):



(II)

wherein:

R<sub>21</sub> is cyano, alkyl, haloalkyl, , acetyl or -C(=S)NH<sub>2</sub>,  
C(=NOH)NH<sub>2</sub> or C(=NNH<sub>2</sub>)NH<sub>2</sub>;

20 R<sub>22</sub> is S(O)<sub>m</sub>R<sub>23</sub>;

R<sub>23</sub> is alkyl or haloalkyl;

R<sub>24</sub> is -N=C(R<sub>25</sub>)N(R<sub>26</sub>)(R<sub>27</sub>) or -N=C(R<sub>25</sub>)-N(R<sub>27</sub>)-R<sub>28</sub>;

R<sub>25</sub> represents hydrogen or alkyl; or alkyl substituted by one or more halogen, alkoxy, haloalkoxy or -S(O)<sub>m</sub>R<sub>35</sub>;

5 R<sub>26</sub> and R<sub>27</sub> each independently represent hydrogen, alkyl, alkenyl or alkynyl; or alkyl substituted by one or more halogen, alkoxy, haloalkoxy, amino, alkylamino, dialkylamino, cyano or -S(O)<sub>m</sub>R<sub>35</sub>; or alkyl substituted by phenyl or pyridyl which rings are optionally substituted with one or more groups selected from halogen, nitro and alkyl; wherein R<sub>35</sub> is alkyl or haloalkyl and m is zero, one or two;

10 X is selected from nitrogen and C-R<sub>32</sub>;

R<sub>28</sub> is alkoxy, haloalkoxy, amino, alkylamino, dialkylamino or -S(O)<sub>t</sub>R<sub>30</sub>;

R<sub>30</sub> is alkyl or haloalkyl;

15 R<sub>31</sub> and R<sub>32</sub> are independently selected from halogen, hydrogen, CN and NO<sub>2</sub>;

R<sub>33</sub> is selected from halogen, haloalkyl, haloalkoxy, -S(O)<sub>r</sub>CF<sub>3</sub>, and -SF<sub>5</sub>;

20 m and r are independently selected from 0, 1, and 2; and t is 0 or 2; with the exclusion of the compound wherein R<sub>21</sub> is cyano; R<sub>22</sub> is -SCF<sub>2</sub>CH<sub>3</sub>; R<sub>25</sub> is hydrogen; X is C-R<sub>32</sub>; R<sub>26</sub> and R<sub>27</sub> are methyl; R<sub>31</sub> and R<sub>32</sub> are chlorine; and R<sub>33</sub> is trifluoromethyl; and veterinarily acceptable salts thereof.

25 6. A compound of formula (XX) or a salt thereof as defined in Claim 2, provided that the compound is not 3-cyano-1-(2,6-dichloro-4-trifluoromethylphenyl)-5-(N-ethoxycarbonyl-N-methyl)amino-4-trifluoromethylthiopyrazole.



5           7.     A composition comprising a parasitically effective,  
substantially non-emetic amount of a compound of formula (I) or a salt  
thereof as defined in Claim 1 or a compound of formula (XX) or a salt  
thereof as defined in Claim 2 and an acceptable carrier.

          8.     The method according to any one of Claims 1 to 4 wherein  
the animal is a domestic animal, preferably a dog or a cat.

10           9.     The method according to any one of Claims 1, 2, 3, 4 or 8  
wherein the 1-arylpyrazole is administered orally in a dose to the animal  
in a dose range generally from 0.1 to 500 mg/kg.

15           10.    The method according to any one of Claims 1, 2, 3, 4, 8, or  
9 wherein the 1-arylpyrazole is administered in a frequency of treatment  
from about once per week to about once per year.

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 99/10452

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D231/44 C07D401/04 A01N43/56

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 301 338 A (BAYER AG) 1 February 1989 (1989-02-01) page 16, line 19 - line 38 ---	1,3,7-10
X	EP 0 811 615 A (RHONE-POULENC AGROCHIMIE) 10 December 1997 (1997-12-10) page 15, line 32 - line 33; example 2 ---	2,6-10
X	EP 0 295 117 A (MAY & BAKER LTD) 14 December 1988 (1988-12-14) page 15, line 9; example 19 page 4, line 47 --- -/-	2,6

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

## \* Special categories of cited documents:

\*A\* document defining the general state of the art which is not considered to be of particular relevance

\*E\* earlier document but published on or after the international filing date

\*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

\*O\* document referring to an oral disclosure, use, exhibition or other means

\*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*Z\* document member of the same patent family

Date of the actual completion of the international search

3 May 2000

Date of mailing of the international search report

15/05/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

De Jong, B

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 99/10452

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE WPI  Section Ch, Week 199818  Derwent Publications Ltd., London, GB;  Class C02, AN 1998-196351  XP002136814  -&amp; JP 10 007509 A (RHONE-POULENC  AGROCHIMIE), 13 January 1998 (1998-01-13)  abstract</p>	2,6-10
X	<p>WO 98 28277 A (RHONE POULENC AGROCHIMIE  ;PHILLIPS JENNIFER (US); PILATO MICHAEL ( )  2 July 1998 (1998-07-02)  page 21, line 9; example 67</p>	1,7-10
X	<p>EP 0 659 745 A (BAYER AG)  28 June 1995 (1995-06-28)  claim 1; example 29</p>	5
X	<p>EP 0 296 381 A (BAYER AG)  28 December 1988 (1988-12-28)  page 6, line 5; claim 15</p>	6
A	<p>WO 98 40359 A (RHONE POULENC AGROCHIMIE  ;WU TAI TEH (US); MANNING DAVID TREADWAY)  17 September 1998 (1998-09-17)  abstract; claim 1</p>	1-10
A	<p>US 5 814 652 A (WU TAI-TEH)  29 September 1998 (1998-09-29)  reference example 1  column 17, line 25 - line 58</p>	1
A	<p>WO 98 28278 A (RHONE POULENC AGROCHIMIE)  2 July 1998 (1998-07-02)  reference examples 21,22  claim 1</p>	1-10

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 99/10452

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0301338 A	01-02-1989	DE 3724919 A	09-02-1989
		AU 603915 B	29-11-1990
		AU 2006288 A	02-02-1989
		BR 8803735 A	14-02-1989
		CA 1335997 A	20-06-1995
		CN 1030912 A,B	08-02-1989
		DD 285279 A	12-12-1990
		DK 96394 A	19-08-1994
		DK 420488 A	29-01-1989
		HU 49045 A,B	28-08-1989
		IL 87210 A	14-01-1993
		JP 1040464 A	10-02-1989
		JP 2634638 B	30-07-1997
		KR 9610342 B	30-07-1996
		MX 171724 B	11-11-1993
		NZ 225540 A	26-04-1990
		PH 25650 A	21-08-1991
		US 4863937 A	05-09-1989
		ZA 8805480 A	26-04-1989
EP 0811615 A	10-12-1997	NONE	
EP 0295117 A	14-12-1988	AT 191479 T	15-04-2000
		AU 618266 B	19-12-1991
		AU 1755488 A	15-12-1988
		BR 8803258 A	31-01-1989
		CA 1330089 A	07-06-1994
		CN 1027341 B	11-01-1995
		CZ 8804052 A	16-12-1998
		DD 281744 B	20-02-1997
		DD 281744 A	22-08-1990
		DK 314088 A	13-12-1988
		EG 19113 A	30-11-1994
		EP 0967206 A	29-12-1999
		FI 882735 A	13-12-1988
		FI 951839 A	18-04-1995
		HU 210668 B	28-06-1995
		HU 48875 A,B	28-07-1989
		HU 9500470 A	30-10-1995
		IL 86492 A	08-07-1993
		IL 105138 A	26-08-1994
		JP 2669538 B	29-10-1997
		JP 63316771 A	26-12-1988
		KR 9701475 B	06-02-1997
		MX 11842 A	01-12-1993
		NO 175367 B	27-06-1994
		NZ 224979 A	26-05-1992
		NZ 236896 A	26-05-1992
		OA 8880 A	31-10-1989
		PH 26895 A	03-12-1992
		PL 272998 A	06-03-1989
		PT 87697 A,B	31-05-1989
		RO 106496 A	31-05-1993
		SG 63529 A	30-03-1999
		SK 405288 A	06-05-1998
		RU 2051909 C	10-01-1996
		TR 23696 A	15-06-1990
		US 5547974 A	20-08-1996

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 99/10452

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0295117 A		US 5714191 A	03-02-1998
		US 5608077 A	04-03-1997
		US 5232940 A	03-08-1993
		US 5916618 A	29-06-1999
		ZA 8804179 A	22-02-1989
		ZW 7388 A	21-12-1988
JP 10007509 A	13-01-1998	NONE	
WO 9828277 A	02-07-1998	AU 5857598 A	17-07-1998
		CZ 9902298 A	15-09-1999
		EP 0948486 A	13-10-1999
		HR 970703 A	31-10-1998
		NO 993107 A	24-08-1999
		PL 334236 A	14-02-2000
EP 0659745 A	28-06-1995	DE 4343832 A	29-06-1995
		AU 6065598 A	18-06-1998
		AU 688401 B	12-03-1998
		AU 8043894 A	06-07-1995
		BR 9405226 A	17-10-1995
		CA 2138452 A	23-06-1995
		CN 1107842 A	06-09-1995
		HU 71745 A	29-01-1996
		JP 8041031 A	13-02-1996
		NZ 270191 A	26-04-1996
		US 5580843 A	03-12-1996
		ZA 9410182 A	25-08-1995
EP 0296381 A	28-12-1988	DE 3719732 A	05-01-1989
		AU 1764488 A	15-12-1988
		BR 8802846 A	03-01-1989
		HU 48217 A	29-05-1989
		JP 63310871 A	19-12-1988
		US 4931461 A	05-06-1990
		ZA 8804152 A	29-03-1989
WO 9840359 A	17-09-1998	AU 6826998 A	29-09-1998
		EP 0973747 A	26-01-2000
		NO 994354 A	04-11-1999
		ZA 9801776 A	03-09-1998
US 5814652 A	29-09-1998	US 5883112 A	16-03-1999
		US 6015910 A	18-01-2000
		AT 223596 A	15-12-1999
		AU 713216 B	25-11-1999
		AU 1302497 A	14-07-1997
		BG 102577 A	30-04-1999
		BR 9612247 A	13-07-1999
		CA 2239081 A	26-06-1997
		CZ 9801932 A	16-09-1998
		DE 19653417 A	26-06-1997
		DK 143796 A	21-06-1997
		WO 9722593 A	26-06-1997
		ES 2124668 A	01-02-1999
		FI 965115 A	21-06-1997
		FR 2742747 A	27-06-1997
		GB 2308365 A, B	25-06-1997

# INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/EP 99/10452

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 5814652 A		GR 96100438 A	29-08-1997
		HR 960596 A	28-02-1998
		IE 960912 A	02-07-1997
		IT MI962693 A	22-06-1998
		JP 2000502095 T	22-02-2000
		LU 88858 A	24-06-1997
		NL 1004863 C	20-10-1998
		NL 1004863 A	02-07-1997
		NO 982802 A	20-08-1998
		PL 327282 A	07-12-1998
		PT 101950 A, B	29-08-1997
		SE 9604636 A	21-06-1997
		SI 9600373 A	31-10-1997
		SK 84498 A	04-11-1998
WO 9828278 A	02-07-1998	AU 5759998 A	17-07-1998
		CZ 9902283 A	17-11-1999
		EP 0948485 A	13-10-1999
		HR 970704 A	31-10-1998
		PL 334282 A	14-02-2000